

INVERSE SPECTRAL PROBLEMS FOR DIRAC OPERATORS WITH SUMMABLE POTENTIALS

S. ALBEVERIO, R. HRYNIV, AND YA. MYKYTYUK

Dedicated to B. M. Levitan, one of the pioneers of this subject

ABSTRACT. The spectral properties of Dirac operators on $(0, 1)$ with potentials that belong entrywise to $L_p(0, 1)$, for some $p \in [1, \infty)$, are studied. The algorithm of reconstruction of the potential from two spectra or from one spectrum and the corresponding norming constants is established, and a complete solution of the inverse spectral problem is provided.

1. INTRODUCTION

The main aim of the present article is to solve the direct and inverse spectral problems for one-dimensional Dirac operators on a finite interval under possibly least restrictive assumptions on their potentials. Namely, the Dirac operators under consideration are generated by the differential expressions

$$\ell_Q := B \frac{d}{dx} + Q(x)$$

and some boundary conditions, where

$$(1.1) \quad B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad Q(x) = \begin{pmatrix} q_1(x) & q_2(x) \\ q_2(x) & -q_1(x) \end{pmatrix},$$

and q_1 and q_2 are real-valued functions from $L_p(0, 1)$, $p \in [1, \infty)$. To simplify unessential technicalities, we shall only consider the boundary conditions that correspond to the Neumann–Dirichlet and Neumann ones in the case of Sturm–Liouville equations, although other boundary conditions can be treated in a similar manner (cf. the study of Sturm–Liouville operators with nonsmooth potentials and various boundary conditions in [26, 52]). The corresponding Dirac operators $\mathcal{A}_1$ and $\mathcal{A}_2$ in the Hilbert space $\mathbb{H} := L_2(0, 1) \times L_2(0, 1)$ act according to the formula $\mathcal{A}_j \mathbf{u} = \ell_Q \mathbf{u}$ on the domains

$$\text{dom } \mathcal{A}_j := \{ \mathbf{u} = (u_1, u_2)^t \mid u_1, u_2 \in \text{AC}(0, 1), \ell_Q \mathbf{u} \in \mathbb{H}, u_2(0) = u_j(1) = 0 \}.$$

It is well known [36] that the operators $\mathcal{A}_1$ and $\mathcal{A}_2$ are selfadjoint in $\mathbb{H}$ and have simple discrete spectra accumulating at $-\infty$ and $+\infty$. Our primary goal is two-fold: firstly, to give a complete description of the spectra of $\mathcal{A}_1$ and $\mathcal{A}_2$ for potentials Q of the form (1.1) with $q_1, q_2 \in L_p(0, 1)$ for some $p \in [1, \infty)$ —i.e., to solve the direct spectral problem,—and, secondly, to give an algorithm of reconstruction of these operators from their spectra or from one spectrum and the corresponding norming constants—i.e., to solve the inverse spectral problem.

Date: February 2, 2008.

2000 Mathematics Subject Classification. Primary 34A55, Secondary 34B30, 47E05.

Key words and phrases. Inverse spectral problems, Dirac operators, non-smooth potentials.

Ever since P. Dirac suggested in 1929 the equation (later named after him) modelling the evolution of spin- $\frac{1}{2}$ particles in the relativistic quantum mechanics [54], its range of applicability in various areas of physics and mathematics has been continuously expanding. In particular, in 1973 Ablowitz, Kaup, Newell, and Segur [1] discovered that the Dirac equation is related to a nonlinear wave equation (the “modified Korteweg–de Vries equation”, a member of the AKNS–ZS hierarchy, see [2, 56]) in the same manner as the Schrödinger equation is related to the KdV equations, and this stimulated the increasing interest in direct and inverse problems for Dirac operators in both physical and mathematical literature. Earlier in 1966, Gasymov and Levitan solved the inverse problems for Dirac operators on $\mathbb{R}_+$ by using the spectral function [16] and by the scattering phase [15]. Their investigations were continued and further developed in many directions. The reference list is so vast, that we can only mention those papers, which, in our opinion, are most pertinent to our topic and refer the reader to the bibliography cited therein for further material. The books by Levitan and Sargsjan [36] and by Thaller [54] may serve as a good introduction to the (respectively mathematical and physical part of the) theory of Dirac operators.

The inverse scattering theory was developed for Dirac operators on the axis in [11, 12, 21, 23, 48], for Dirac systems of order $2n$ on semiaxis in [13], and for more general canonical systems on $\mathbb{R}$ in [50]. The nonselfadjoint case was treated in [37] and nonstationary scattering, including point interactions, in [47] and [4], respectively. Reconstruction from the spectral function on semiaxis was done in [51] for a general boundary condition at $x = 0$ and in [53] in the case of an interface condition in an interior point; the general first order systems in $L_2(\mathbb{R}_+, \mathbb{C}^{2n})$ were recently treated in [33]. Inverse problems in the periodic case were studied in [29, 31], and the Weyl–Titchmarsh m -function was used to recover the potential of the Dirac operator in [5] and of the Dirac systems of order $2n$ in [7, 17, 49] (see also the detailed reference lists therein).

The inverse problems for Dirac operators on a finite interval have also been studied in detail. Reconstruction of a continuous potential from two spectra was carried out in [14], from one spectrum and the norming constants (in the presence of a Coulomb-type singularity) in [9], and from the spectral function in [40]. Explicit formulae for solutions (based on the degenerate Gelfand–Levitan–Marchenko equation) in the case where finitely many spectral data are perturbed were given in [8]. Uniqueness results for other types of inverse problems were established—e.g., for mixed spectral [25] or interior [42] data, nonseparated boundary conditions [45], or for the weighted Dirac equations [55]. Ambartsumyan-type theorems were proved in [24] and for the matrix case in [30]. Finally, uniqueness of the inverse problem for general Dirac-type systems of order $2n$ was recently established in [38, 39].

We observe that in the above-cited papers the inverse spectral problems for Dirac operators on a finite interval were considered for continuous potentials only, which excludes, e.g., the important case of piecewise constant potentials. We remove this restriction by allowing potentials belonging entrywise to $L_p(0, 1)$, $p \in [1, \infty)$, and completely solve the inverse spectral problem for Dirac operators in this class (see Theorems 3.1, 3.3, 5.1, and 5.7). The main idea of the proof rests on the fact that the transformation operators for Dirac operators under consideration satisfy not only the classical Gelfand–Levitan–Marchenko equation (4.4), but also its counterpart (4.9), which was used by Krein [32] in the study of the inverse problem for impedance Sturm–Liouville equations (see also [3, 10]). This “Krein equation” survives the passage to

the limit in the L_p -topology and thus allows us to treat potentials belonging to $L_p(0, 1)$ entrywise.

The paper is organized as follows. In Section 2 transformation operators are constructed and some of their properties are established. Based on this, in Section 3 we find the asymptotics of eigenvalues and norming constants for the operators $\mathcal{A}_1$ and $\mathcal{A}_2$. The Gelfand–Levitan–Marchenko and Krein equations, which relate the spectral data and the transformation operators, are derived in Section 4, and the solution of the inverse spectral problem is given in Section 5. Finally, two appendices contain some facts related to harmonic analysis and the factorisation theory in operator algebras.

Throughout the paper, we shall denote by $\langle \cdot, \cdot \rangle$ the scalar product in $\mathbb{H}$ and by $\mathcal{M}_2$ the algebra of 2×2 matrices with complex entries endowed with the operator norm $|\cdot|$ of the Euclidean space $\mathbb{C}^2$. Where no confusion arises, we abbreviate $L_p(0, 1)$ to L_p and write $L_p(\mathcal{M}_2)$ for the space $L_p((0, 1), \mathcal{M}_2)$ of $\mathcal{M}_2$ -valued functions on $(0, 1)$ with complex-valued entries and the norm

$$\|V\|_{L_p} := \left(\int_0^1 |V(t)|^p dt \right)^{1/p}.$$

Also, $(x, y)^t$ shall stand for the column-vector in $\mathbb{C}^2$ with components x and y .

2. TRANSFORMATION OPERATORS

Assume that $Q \in L_p(\mathcal{M}_2)$, $p \in [1, \infty)$, is of the form (1.1) and denote by $U(\cdot) = U(\cdot, \lambda)$ the Cauchy matrix corresponding to the equation $\ell_Q \mathbf{u} = \lambda \mathbf{u}$. In other words, U is a 2×2 matrix-valued function satisfying the equation

$$(2.1) \quad B \frac{dU}{dx} + QU = \lambda U$$

and the initial condition $U(0) = I := \text{diag}(1, 1)$. Denoting by $\mathbf{c}(\cdot, \lambda) := (c_1(\cdot, \lambda), c_2(\cdot, \lambda))^t$ and $\mathbf{s}(\cdot, \lambda) := (s_1(\cdot, \lambda), s_2(\cdot, \lambda))^t$ the solutions of the equation $\ell_Q \mathbf{u} = \lambda \mathbf{u}$ satisfying the initial conditions $c_1(0, \lambda) = s_2(0, \lambda) = 1$ and $c_2(0, \lambda) = s_1(0, \lambda) = 0$, we find that

$$U(x, \lambda) = \begin{pmatrix} c_1(x, \lambda) & s_1(x, \lambda) \\ c_2(x, \lambda) & s_2(x, \lambda) \end{pmatrix}.$$

Our next aim is to derive an integral representation for U of a special form.

Theorem 2.1. *Assume that $Q \in L_p(\mathcal{M}_2)$, $p \in [1, \infty)$. Then*

$$(2.2) \quad U(x, \lambda) = e^{-\lambda x B} + \int_0^x e^{-\lambda(x-2s)B} P(x, s) ds,$$

where the matrix-valued function $P = P_Q$ has the following properties:

- (a) for every $x \in [0, 1]$ the function $P(x, \cdot)$ belongs to $L_p(\mathcal{M}_2)$;
- (b) the mapping $\mathcal{P}_Q : x \mapsto P_Q(x, \cdot) \in L_p(\mathcal{M}_2)$ is continuous on $[0, 1]$;
- (c) the function $\mathcal{P}_Q$ depends continuously in $C([0, 1], L_p(\mathcal{M}_2))$ on $Q \in L_p(\mathcal{M}_2)$.

Proof. The standard variation of constant arguments show that U satisfies the equivalent integral equation (recall that $B^2 = -I$)

$$(2.3) \quad U(x, \lambda) = e^{-\lambda x B} + \int_0^x e^{-\lambda(x-t)B} BQ(t)U(t) dt,$$

which can be solved by the method of successive approximations. Namely, with

$$(2.4) \quad U_0(x) := e^{-\lambda x B} \quad \text{and} \quad U_{n+1}(x) := \int_0^x e^{-\lambda(x-t)B} BQ(t)U_n(t) dt \quad \text{for } n \geq 0,$$

the solution of (2.3) formally equals $\sum_{n=0}^{\infty} U_n$. Assume that we have proved that

$$(2.5) \quad \sum_{n=0}^{\infty} \|U_n\|_{\infty} < \infty,$$

where $\|U_n\|_{\infty} := \sup_{x \in [0,1]} |U_n(x)|$. Differentiating then the recurrence relations (2.4), we find that

$$U'_n(x) = -\lambda B U_n(x) + BQ(x)U_{n-1}(x),$$

which in view of (2.5) shows that the series $\sum_{n=0}^{\infty} U_n$ converges in the topology of the space $W_p^1((0,1), \mathcal{M}_2)$ to some $\mathcal{M}_2$ -valued function V . This function V solves (2.1) and satisfies the initial condition $V(0) = I$, and hence it coincides with the Cauchy matrix U .

To justify (2.5), we use the identity

$$(2.6) \quad e^{-\lambda x B} Q(t) = Q(t) e^{\lambda x B}, \quad x, t \in [0,1],$$

in the recurrence relations (2.4) and derive the formula

$$(2.7) \quad U_n(x) = \int_{\Pi_n(x)} e^{-\lambda(x-2\xi_n(t))B} BQ(t_1) \cdots BQ(t_n) dt_1 \cdots dt_n,$$

in which we have set

$$\begin{aligned} \Pi_n(x) &= \{t := (t_1, \dots, t_n) \in \mathbb{R}^n \mid 0 \leq t_n \leq \cdots \leq t_1 \leq x\}, \\ \xi_n(t) &= \sum_{l=1}^n (-1)^{l+1} t_l. \end{aligned}$$

Upon the change of variables $s = \xi_n(t)$, $y_l = t_{l+1}$, $l = 1, 2, \dots, n-1$, we recast the integral in (2.7) as

$$U_n(x) = \int_0^x e^{-\lambda(x-2s)B} P_n(x, s) ds,$$

where $P_1(x, s) \equiv BQ(s)$ and, for all $n \in \mathbb{N}$ and $0 \leq s \leq x \leq 1$,

$$(2.8) \quad P_{n+1}(x, s) = \int_{\Pi_n^*(x, s)} BQ(s + \xi_n(y)) BQ(y_1) \cdots BQ(y_n) dy_1 \cdots dy_n,$$

with

$$\Pi_n^*(x, s) = \{y = (y_1, \dots, y_n) \in \mathbb{R}^n \mid 0 \leq y_n \leq y_{n-1} \leq \cdots \leq y_1 \leq s + \xi_n(y) \leq x\}.$$

For convenience, we extend the functions P_n , $n \geq 2$, to the whole square $[0,1] \times [0,1]$ by setting $P_n(x, s) = 0$ for $0 \leq x < s \leq 1$.

Using the Hölder inequality and Fubini's theorem, we find that, for every $n \in \mathbb{N}$, the function $P_{n+1}(x, \cdot)$ belongs to $L_p(\mathcal{M}_2)$ and that

$$\begin{aligned} \|P_{n+1}(x, \cdot)\|_{L_p}^p &= \int_0^1 |P_{n+1}(x, s)|^p ds \\ &\leq (n!)^{1-p} \int_0^1 \int_{\Pi_n^*(x, s)} |Q(s + \xi_{n-1}(y))|^p |Q(y_1)|^p \cdots |Q(y_n)|^p dy_1 \dots dy_n ds \\ &= (n!)^{1-p} \int_{\Pi_{n+1}(x)} |Q(t_1)|^p \cdots |Q(t_{n+1})|^p dt_1 \dots dt_{n+1} \\ &= \frac{1}{(n!)^p (n+1)} \left(\int_0^x |Q|^p \right)^{n+1} \leq \frac{\|Q\|_{L_p}^{(n+1)p}}{(n!)^p}. \end{aligned}$$

Henceforth with $C := \max_{x \in [-1, 1]} |e^{-\lambda x B}|$ we have

$$|U_n(x)| \leq C \int_0^x |P_n(x, s)| ds \leq C \|P_n(x, \cdot)\|_{L_p} \leq C \frac{\|Q\|_{L_p}^n}{(n-1)!},$$

and (2.5) follows.

Moreover, the above inequality implies that the series $\sum_{n=1}^{\infty} P_n(x, \cdot)$ converges in $L_p(\mathcal{M}_2)$ to some function $P(x, \cdot)$ and yields the estimate

$$(2.9) \quad \|P(x, \cdot)\|_{L_p} \leq \sum_{n=1}^{\infty} \frac{\|Q\|_{L_p}^n}{(n-1)!} = \|Q\|_{L_p} \exp\{\|Q\|_{L_p}\}$$

for all $x \in [0, 1]$. This establishes (a).

Assume that $\tilde{Q}$ is another potential in $L_p(\mathcal{M}_2)$ and denote by $\tilde{P}_n$ the corresponding functions constructed as above but for $\tilde{Q}$ instead of Q ; then similar calculations on account of the inequality

$$\begin{aligned} \left| \prod_{k=1}^n a_k - \prod_{k=1}^n b_k \right|^p &\leq \left(\sum_{k=1}^n |a_k - b_k| \prod_{j \neq k} (|a_j| + |b_j|) \right)^p \\ &\leq n^{p-1} \sum_{k=1}^n |a_k - b_k|^p \prod_{j \neq k} (|a_j| + |b_j|)^p \end{aligned}$$

lead to the estimate

$$(2.10) \quad \|P_{n+1}(x, \cdot) - \tilde{P}_{n+1}(x, \cdot)\|_{L_p}^p \leq \left(\frac{n+1}{n!} \right)^p \|Q - \tilde{Q}\|_{L_p}^p \left(\|Q\|_{L_p} + \|\tilde{Q}\|_{L_p} \right)^{np}.$$

It follows that

$$(2.11) \quad \|P_Q(x, \cdot) - P_{\tilde{Q}}(x, \cdot)\|_{L_p} \leq (1 + 2r) e^{2r} \|Q - \tilde{Q}\|_{L_p}$$

as soon as r is such that $\|Q\|_{L_p}, \|\tilde{Q}\|_{L_p} \leq r$.

Observe that if $\tilde{Q} \in C([0, 1], \mathcal{M}_2)$, then the functions $\tilde{P}_n$, $n \geq 2$, are continuous in the square $[0, 1] \times [0, 1]$, and, moreover,

$$\max_{0 \leq x, s \leq 1} |\tilde{P}_n(x, s)| \leq \frac{\|\tilde{Q}\|_{\infty}^n}{n!},$$

so that the function $[0, 1] \ni x \mapsto P_{\tilde{Q}}(x, \cdot) \in L_p(\mathcal{M}_2)$ is continuous. Since the potential $Q \in L_p(\mathcal{M}_2)$ is the limit in $L_p(\mathcal{M}_2)$ of potentials $\tilde{Q}_n \in C([0, 1], \mathcal{M}_2)$, estimate (2.11) yields both assertions (b) and (c). The proof is complete. $\square$

Corollary 2.2. *Assume that $Q \in L_p(\mathcal{M}_2)$ and set*

$$P^+ = \sum_{n=1}^{\infty} P_{2n}, \quad P^- = \sum_{n=1}^{\infty} P_{2n-1},$$

where the functions P_n are given by formula (2.8). Set $\mathbf{c}_0(x, \lambda) := (\cos \lambda x, \sin \lambda x)^t$ and

$$(2.12) \quad \begin{aligned} R(x, t) &= R_Q(x, t) := P^+(x, t) + P^-(x, t)J, \\ K(x, t) &= K_Q(x, t) := \frac{1}{2} \left[R(x, \frac{x-t}{2}) + R(x, \frac{x+t}{2})J \right], \end{aligned}$$

where $J = \text{diag}\{1, -1\}$. Then the vector-function $\mathbf{c}(\cdot, \lambda)$ is given by

$$(2.13) \quad \mathbf{c}(x, \lambda) = \mathbf{c}_0(x, \lambda) + \int_0^x K(x, t) \mathbf{c}_0(t, \lambda) dt.$$

Proof. Using (2.6) and (2.8), we conclude that

$$\begin{aligned} e^{-\lambda(x-2s)B} P_{2n}(x, s) &= P_{2n}(x, s) e^{-\lambda(x-2s)B}, \\ e^{-\lambda(x-2s)B} P_{2n-1}(x, s) &= P_{2n-1}(x, s) e^{\lambda(x-2s)B}. \end{aligned}$$

Therefore equality (2.2) can be written as

$$U(x, \lambda) = e^{-\lambda x B} + \int_0^x P^+(x, s) e^{-\lambda(x-2s)B} ds + \int_0^x P^-(x, s) e^{\lambda(x-2s)B} ds.$$

Observing that

$$e^{-\lambda x B} = \begin{pmatrix} \cos \lambda x & -\sin \lambda x \\ \sin \lambda x & \cos \lambda x \end{pmatrix}$$

and taking the first column of the above equality, we get

$$\mathbf{c}(x, \lambda) = \mathbf{c}_0(x, \lambda) + \int_0^x R(x, s) \mathbf{c}_0(x - 2s, \lambda) ds.$$

Since

$$\begin{aligned} \int_0^{x/2} R(x, s) \mathbf{c}_0(x - 2s, \lambda) ds &= \frac{1}{2} \int_0^x R(x, \frac{x-t}{2}) \mathbf{c}_0(t, \lambda) dt \\ \int_{x/2}^x R(x, s) \mathbf{c}_0(x - 2s, \lambda) ds &= \int_{x/2}^x R(x, s) J \mathbf{c}_0(2s - x, \lambda) ds \\ &= \frac{1}{2} \int_0^x R(x, \frac{x+t}{2}) J \mathbf{c}_0(t, \lambda) dt, \end{aligned}$$

the required relation follows. $\square$

Equality (2.13) shows that the operator $\mathcal{J} + \mathcal{K}$ defined by

$$(\mathcal{J} + \mathcal{K})\mathbf{u}(x) = \mathbf{u}(x) + \int_0^x K(x, t) \mathbf{u}(t) dt$$

transforms the solution of the equation $\ell_0 \mathbf{u} = \lambda \mathbf{u}$ (i.e., with a potential Q equal to zero identically) subject to the initial conditions $u_1(0) = 1, u_2(0) = 0$ into the solution

of the equation $\ell_Q \mathbf{u} = \lambda \mathbf{u}$ satisfying the same initial conditions. Denote by $\tilde{\mathcal{A}}_Q$ the operator in $\mathbb{H}$ acting as $\tilde{\mathcal{A}}_Q \mathbf{u} = \ell_Q \mathbf{u}$ on the domain

$$\text{dom}(\tilde{\mathcal{A}}_Q) := \{\mathbf{u} = (u_1, u_2)^t \in W_2^1(0, 1) \times W_2^1(0, 1) \mid u_2(0) = 0\};$$

then $\mathcal{I} + \mathcal{K}$ is in fact the transformation operator for $\tilde{\mathcal{A}}_Q$ and $\tilde{\mathcal{A}}_0$, i.e., $\tilde{\mathcal{A}}_Q(\mathcal{I} + \mathcal{K}) = (\mathcal{I} + \mathcal{K})\tilde{\mathcal{A}}_0$, see Theorem 2.4.

The operator $\mathcal{K}$ possesses some important properties, which we now establish. Denote by $G_p(\mathcal{M}_2)$ the set of measurable 2×2 matrix-valued functions K on $[0, 1] \times [0, 1]$ having the property that, for each x and t in $[0, 1]$, the matrix-valued functions $K(x, \cdot)$ and $K(\cdot, t)$ belong to $L_p(\mathcal{M}_2)$ and, moreover, the mappings

$$[0, 1] \ni x \mapsto K(x, \cdot) \in L_p(\mathcal{M}_2), \quad [0, 1] \ni t \mapsto K(\cdot, t) \in L_p(\mathcal{M}_2)$$

are continuous (i.e., they coincide a.e. with some continuous mappings from $[0, 1]$ into $L_p(\mathcal{M}_2)$). The set $G_p(\mathcal{M}_2)$ becomes a Banach space under the norm

$$(2.14) \quad \|K\|_{G_p} := \max \left\{ \max_{x \in [0, 1]} \|K(x, \cdot)\|_{L_p}, \max_{t \in [0, 1]} \|K(\cdot, t)\|_{L_p} \right\}.$$

We also denote by $\mathcal{G}_p(\mathcal{M}_2)$ the set of the integral operators $\mathcal{K}$ in $\mathbb{H}$ with kernels K from $G_p(\mathcal{M}_2)$. Under the induced norm $\|\mathcal{K}\|_{\mathcal{G}_p} := \|K\|_{G_p}$, the set $\mathcal{G}_p(\mathcal{M}_2)$ becomes an algebra. The algebra $\mathcal{G}_p(\mathcal{M}_2)$ is continuously embedded into the algebra $\mathcal{B}(\mathbb{H})$ of all bounded operators in $\mathbb{H}$ since the functions K belonging to $G_p(\mathcal{M}_2)$ have finite Holmgren norm [22]; moreover, for $\mathcal{K} \in \mathcal{G}_p(\mathcal{M}_2)$ the inequality $\|\mathcal{K}\|_{\mathcal{B}(\mathbb{H})} \leq \|\mathcal{K}\|_{\mathcal{G}_p}$ holds true.

Theorem 2.3. *Assume that $Q \in L_p(\mathcal{M}_2)$; then the integral operator $\mathcal{K} = \mathcal{K}_Q$ with kernel K of (2.12) belongs to $\mathcal{G}_p(\mathcal{M}_2)$ and, moreover, the mapping $L_p(\mathcal{M}_2) \ni Q \mapsto \mathcal{K}_Q \in \mathcal{G}_p(\mathcal{M}_2)$ is continuous.*

Proof. In view of relations (2.12), we have

$$(2.15) \quad K(x, t) = \frac{1}{2} \left[P^+(x, \frac{x-t}{2}) + P^-(x, \frac{x-t}{2})J + P^+(x, \frac{x+t}{2})J + P^-(x, \frac{x+t}{2}) \right],$$

and hence it suffices to prove the assertions of the theorem for the operators with kernels $P^+(x, \frac{x \pm t}{2})$ and $P^-(x, \frac{x \pm t}{2})$. Since the proof is analogous for all four functions, we shall give it for the function $P^+(x, \frac{x-t}{2}) =: \hat{P}(x, t)$ only.

It follows from the proof of Theorem 2.1 that the function P^+ enjoys the properties (a)–(c) of that theorem. Simple arguments based on the change of variables justify the validity of the properties (a)–(c) for the kernel $\hat{P}$. It thus remains to establish similar properties of $\hat{P}$ with respect to the variable t .

Assume first that Q is continuous. Changing the variables $\eta = s + \xi(y)$, $\tilde{y}_1 = y_2, \dots, \tilde{y}_{n-1} = y_n$ in integral (2.8), we arrive at the relation

$$P_{n+1}(x, s) = \int_s^x BQ(\eta)P_n(\eta, \eta - s) d\eta,$$

which yields

$$P^+(x, s) = \int_s^x BQ(\eta)P^-(\eta, \eta - s) d\eta.$$

It follows that

$$\begin{aligned} \|\hat{P}(\cdot, t)\|_{L_p}^p &= \int_t^1 \left| \int_{\frac{x-t}{2}}^x BQ(\eta) P^-(\eta, \eta - \frac{x-t}{2}) d\eta \right|^p dx \\ &\leq \int_0^1 d\eta |Q(\eta)|^p \int_t^{2\eta+t} |P^-(\eta, \eta - \frac{x-t}{2})|^p dx \\ &\leq \|Q\|_{L_p}^p \max_{\eta \in [0,1]} \|P^-(\eta, \cdot)\|_{L_p}^p \leq \|Q\|^{2p} \exp\{p\|Q\|_{L_p}\}, \end{aligned}$$

cf. (2.9). If $\tilde{Q}$ is another continuous potential, then we find analogously that

$$\begin{aligned} \|\hat{P}_Q(\cdot, t) - \hat{P}_{\tilde{Q}}(\cdot, t)\|_{L_p}^p &\leq 2^{p-1} \|Q - \tilde{Q}\|_{L_p}^p \max_{\eta \in [0,1]} \|P_Q^-(\eta, \cdot)\|_{L_p}^p \\ &\quad + 2^{p-1} \|\tilde{Q}\|_{L_p}^p \max_{\eta \in [0,1]} \|P_{\tilde{Q}}^-(\eta, \cdot) - P_Q^-(\eta, \cdot)\|_{L_p}^p. \end{aligned}$$

Recalling inequality (2.10), we conclude that the function $t \mapsto \hat{P}_Q(\cdot, t)$, which belongs to $C([0, 1], L_p(\mathcal{M}_2))$ if Q is continuous, depends therein continuously on $Q \in C([0, 1], \mathcal{M}_2)$ with respect to the topology of $L_p(\mathcal{M}_2)$. Since the space $C([0, 1], \mathcal{M}_2)$ is dense in $L_p(\mathcal{M}_2)$, we show by continuity that, for every $Q \in L_p(\mathcal{M}_2)$, the function $\hat{P}_Q(\cdot, t)$ belongs to $L_p(\mathcal{M}_2)$ for every fixed $t \in [0, 1]$, that the mapping $t \mapsto \hat{P}_Q(\cdot, t)$ is continuous, and that the continuous $L_p(\mathcal{M}_2)$ -valued function of t , $t \mapsto P_Q(\cdot, t)$, depends continuously in $C([0, 1], L_p(\mathcal{M}_2))$ on $Q \in L_p(\mathcal{M}_2)$.

Summing up, we have shown that the function $\hat{P} = \hat{P}_Q$ belongs to $G_p(\mathcal{M}_2)$ and depends in $G_p(\mathcal{M}_2)$ continuously on $Q \in L_p(\mathcal{M}_2)$. This establishes the theorem. $\square$

Theorem 2.4. *Assume that $Q \in L_p(\mathcal{M}_2)$ and let $\mathcal{K}$ be an integral operator with kernel K of (2.12). Then $\mathcal{I} + \mathcal{K}$ is the transformation operator for the pair $\tilde{\mathcal{A}}_Q$ and $\tilde{\mathcal{A}}_0$, i.e., $\tilde{\mathcal{A}}_Q(\mathcal{I} + \mathcal{K}) = (\mathcal{I} + \mathcal{K})\tilde{\mathcal{A}}_0$.*

Proof. Since $\mathcal{K}$ belongs to $\mathcal{G}_p(\mathcal{M}_2)$ by Theorem 2.3 and its kernel K is lower-diagonal, it follows that $\mathcal{K}$ is a Volterra operator in $\mathbb{H}$ and hence $\mathcal{I} + \mathcal{K}$ is a homeomorphism of $\mathbb{H}$.

Write $\hat{\mathcal{A}}_Q := (\mathcal{I} + \mathcal{K})^{-1} \tilde{\mathcal{A}}_Q (\mathcal{I} + \mathcal{K})$. In view of (2.13), $\mathbf{c}_0(\cdot, \lambda)$ is an eigenvector of the operator $\hat{\mathcal{A}}_Q$ corresponding to the eigenvalue λ for every $\lambda \in \mathbb{C}$. Denote by $\mathcal{L}$ the linear hull of the system $\{\mathbf{c}_0(\cdot, \lambda) \mid \lambda \in \mathbb{C}\}$; then the restrictions of the operators $\hat{\mathcal{A}}_Q$ and $\tilde{\mathcal{A}}_0$ onto $\mathcal{L}$ coincide. Since $\mathcal{L}$ is a core of $\tilde{\mathcal{A}}_0$ (see a similar result in [3, Theorem 3.3]) and $\hat{\mathcal{A}}_Q$ is closed, it follows that $\tilde{\mathcal{A}}_0 \subset \hat{\mathcal{A}}_Q$. It remains to observe that $\hat{\mathcal{A}}_Q$ cannot be a proper extension of $\tilde{\mathcal{A}}_0$ since otherwise $\hat{\mathcal{A}}_Q$ —and $\tilde{\mathcal{A}}_Q$ by similarity—would have a two-dimensional nullspace, which would contradict the uniqueness of solutions to the equation $\ell_Q \mathbf{u} = \lambda \mathbf{u}$. Thus $\tilde{\mathcal{A}}_0 = \hat{\mathcal{A}}_Q$, and the proof is complete. $\square$

3. DIRECT SPECTRAL PROBLEM

The aim of this section is to perform the direct spectral analysis for the Dirac operators $\mathcal{A}_1$ and $\mathcal{A}_2$. The main tool of our investigations will be the transformation operators constructed in the previous section.

Theorem 3.1. *Assume that $Q \in L_p(\mathcal{M}_2)$; then the eigenvalues $(\lambda_n)_{n \in \mathbb{Z}}$ and $(\mu_n)_{n \in \mathbb{Z}}$ of $\mathcal{A}_1$ and $\mathcal{A}_2$ respectively can be enumerated so that they satisfy the interlacing condition*

$$(3.1) \quad \lambda_{n-1} < \mu_n < \lambda_n, \quad n \in \mathbb{Z},$$

and the asymptotics

$$(3.2) \quad \begin{aligned} \lambda_n &= \pi(n + \tfrac{1}{2}) + e_n(g_1), \\ \mu_n &= \pi n + e_n(g_2), \end{aligned}$$

where $g_1, g_2 \in L_p$ and $e_n(g) := \int_0^1 e^{-2\pi nix} g(x) dx$ are the Fourier coefficients of a function g .

Proof. The fact that the spectra of $\mathcal{A}_1$ and $\mathcal{A}_2$ interlace (i.e., that between two consecutive eigenvalues of one operator, there is exactly one eigenvalue of the other operator) is well known (see, e.g., [14]). It is easily seen that if we have an enumeration of λ_n and μ_n obeying (3.2) for some $g_1, g_2 \in L_p$, then (3.1) holds for all n with sufficiently large $|n|$. Since the two spectra interlace, we can permute a finite number of indices if necessary in such a way that (3.1) becomes valid for all $n \in \mathbb{Z}$. This reordering amounts to adding trigonometric polynomials of finite degree to the functions g_1 and g_2 , and the modified functions g_1 and g_2 will remain in L_p . Therefore it suffices to establish (3.2).

The equation $\ell_Q \mathbf{u} = \lambda \mathbf{u}$ subject to the initial conditions $f_1(0) = 1, f_2(0) = 0$ has the solution

$$\mathbf{c}(x, \lambda) = \mathbf{c}_0(x, \lambda) + \int_0^x K(x, s) \mathbf{c}_0(s, \lambda) ds,$$

where $K = (k_{jl})_{j,l=1}^2$ is the kernel of the transformation operator constructed in Corollary 2.2. The numbers λ_n are zeros of the function $c_1(1, \lambda)$, which, after simple transformations, takes the form

$$(3.3) \quad \begin{aligned} c_1(1, \lambda) &= \cos \lambda + \int_0^1 [k_{11}(s) \cos(\lambda s) + k_{12}(s) \sin(\lambda s)] ds \\ &= \cos \lambda + \int_{-1}^1 f_1(s) e^{i\lambda s} ds, \end{aligned}$$

where

$$f_1(s) := \begin{cases} \frac{1}{2} [k_{11}(1, s) - ik_{12}(1, s)], & s \geq 0, \\ \frac{1}{2} [k_{11}(1, -s) + ik_{12}(1, -s)], & s < 0 \end{cases}$$

is a function in $L_p(-1, 1)$.

Analogously the eigenvalues of $\mathcal{A}_2$ are zeros of the entire function $c_2(1, \lambda)$, which has the form

$$(3.4) \quad c_2(1, \lambda) = \sin \lambda + \int_{-1}^1 f_2(s) e^{i\lambda s} ds$$

for some $f_2 \in L_p(-1, 1)$.

The required asymptotics for zeros of $c_1(1, \lambda)$ and $c_2(1, \lambda)$ follows now from [27]. $\square$

Definition 3.2. We denote by SD_p the set of all pairs $\{(\lambda_n)_{n \in \mathbb{Z}}, (\mu_n)_{n \in \mathbb{Z}}\}$, in which (λ_n) and (μ_n) are sequences of real numbers that obey the interlacing condition (3.1) and the asymptotics (3.2).

Theorem 3.1 shows that, for any real-valued $Q \in L_p(\mathcal{M}_2)$, the spectra of the operators $\mathcal{A}_1$ and $\mathcal{A}_2$ form an element of SD_p . In the reverse direction, Theorem 5.1 claims that any element of SD_p is composed of the spectra of Dirac operators $\mathcal{A}_1$ and $\mathcal{A}_2$ for some $Q \in L_p(\mathcal{M}_2)$. The reconstruction algorithm uses in fact the spectrum (λ_n) of

$\mathcal{A}_1$ and the sequence of corresponding norming constants $\alpha_n := \|\mathbf{c}(\cdot, \lambda_n)\|^{-2}$, whose properties we are going to study next.

Theorem 3.3. *Assume that $Q \in L_p(\mathcal{M}_2)$; then the norming constants $\alpha_n = \|\mathbf{c}(\cdot, \lambda_n)\|^{-2}$ have the asymptotics*

$$(3.5) \quad \alpha_n = 1 + e_n(g),$$

where $g \in L_p$.

Set $\phi(\lambda) := c_1(1, \lambda)$ and $\psi(\lambda) := c_2(1, \lambda)$. We observe that the functions $\phi(\lambda)$ and $\psi(\lambda)$ can be reconstructed from their zeros as follows [27].

Proposition 3.4. *The following equalities hold:*

$$(3.6) \quad \phi(\lambda) = \text{V.p.} \prod_{n=-\infty}^{\infty} \frac{\lambda_n - \lambda}{\pi(n + \frac{1}{2})}, \quad \psi(\lambda) = (\lambda - \mu_0) \text{V.p.} \prod_{n=-\infty}^{\infty} \frac{\mu_n - \lambda}{\pi n}$$

(where the prime means that the factor corresponding to the index $n = 0$ should be omitted); moreover, the products converge uniformly on compact sets.

It turns out that ϕ and ψ (i.e., that the two spectra $(\lambda_n)_{n \in \mathbb{Z}}$ and $(\mu_n)_{n \in \mathbb{Z}}$) determine the norming constants α_n as follows.

Lemma 3.5. *The norming constants α_n satisfy the following relation:*

$$(3.7) \quad \alpha_n = -\frac{1}{\phi(\lambda_n)\psi(\lambda_n)}.$$

Proof. Since the system of eigenvectors $(\mathbf{c}_n)_{n \in \mathbb{Z}}$, $\mathbf{c}_n := \mathbf{c}(\cdot, \lambda_n)$, of the operator $\mathcal{A}_1$ is an orthogonal basis of $\mathbb{H}$, we have

$$(\mathcal{A}_1 - \lambda)^{-1} \mathbf{u} = \sum_{n=-\infty}^{\infty} \frac{\alpha_n \langle \mathbf{u}, \mathbf{c}_n \rangle \mathbf{c}_n}{\lambda_n - \lambda}.$$

In particular, the residue of this expression at $\lambda = \lambda_n$ equals $-\alpha_n \langle \mathbf{u}, \mathbf{c}_n \rangle \mathbf{c}_n$.

On the other hand, $(\mathcal{A}_1 - \lambda)^{-1}$ can be calculated as

$$\begin{aligned} (\mathcal{A}_1 - \lambda)^{-1} \mathbf{u}(x) &= \left[\int_x^1 \langle \mathbf{u}(t), \overline{\mathbf{s}^+(t, \lambda)} \rangle_{\mathbb{C}^2} dt \right] \frac{\mathbf{c}(x, \lambda)}{W(\lambda)} \\ &\quad + \left[\int_0^x \langle \mathbf{u}(t), \overline{\mathbf{c}(t, \lambda)} \rangle_{\mathbb{C}^2} dt \right] \frac{\mathbf{s}^+(x, \lambda)}{W(\lambda)}, \end{aligned}$$

where $\mathbf{s}^+(\cdot, \lambda)$ is a solution of the equation $\ell_Q(\mathbf{u}) = \lambda^2 \mathbf{u}$ subject to the terminal conditions $u_1(1) = 0$ and $u_2(1) = 1$, and $W(\lambda)$ is the Wronskian of the solutions $\mathbf{c}(\cdot, \lambda)$ and $\mathbf{s}^+(\cdot, \lambda)$, i.e.,

$$W(\lambda) := c_1(x, \lambda) s_2^+(x, \lambda) - c_2(x, \lambda) s_1^+(x, \lambda).$$

Since the right-hand side of the above equality does not depend on x , we see that

$$W(\lambda) = c_1(1, \lambda) = \phi(\lambda).$$

Thus the residue of $(\mathcal{A}_1 - \lambda)^{-1} \mathbf{u}$ at $\lambda = \lambda_n$ is equal to

$$(3.8) \quad \left[\int_x^1 \langle \mathbf{u}(t), \overline{\mathbf{s}^+(t, \lambda_n)} \rangle_{\mathbb{C}^2} dt \right] \frac{\mathbf{c}(x, \lambda_n)}{\phi(\lambda_n)} + \left[\int_0^x \langle \mathbf{u}(t), \overline{\mathbf{c}(t, \lambda_n)} \rangle_{\mathbb{C}^2} dt \right] \frac{\mathbf{s}^+(x, \lambda_n)}{\phi(\lambda_n)}.$$

We observe now that the vector-functions $\mathbf{s}^+(x, \lambda_n)$ and $\mathbf{c}(x, \lambda_n)$ are collinear, namely,

$$\mathbf{s}^+(\cdot, \lambda_n) = \frac{s_2^+(1, \lambda_n)}{c_2(1, \lambda_n)} \mathbf{c}(\cdot, \lambda_n) = \frac{1}{\psi(\lambda_n)} \mathbf{c}(\cdot, \lambda_n),$$

and thus expression (3.8) simplifies to $\langle \mathbf{u}, \mathbf{c}_n \rangle \mathbf{c}_n / (\dot{\phi}(\lambda_n) \psi(\lambda_n))$. Equating this with the above expression for the residue of $(\mathcal{A}_1 - \lambda)^{-1} \mathbf{u}$ at $\lambda = \lambda_n$, we obtain relation (3.7) for α_n . The proof is complete. $\square$

Proof of Theorem 3.3. In view of equality (3.7) and Propositions A.1 and A.2 it suffices to prove that the numbers $a_n := -\dot{\phi}(\lambda_n)$ and $b_n := \psi(\lambda_n)$ can be represented as $(-1)^n(1 + \tilde{a}_n)$ and $(-1)^n(1 + \tilde{b}_n)$ respectively, where $\tilde{a}_n$ and $\tilde{b}_n$ are n -th Fourier coefficients of some functions from L_p .

Using formulae (3.3) and (3.4), we show that the numbers a_n and b_n are of the form

$$\sin \lambda_n + \int_{-1}^1 f(s) e^{i\lambda_n s} ds$$

with $f(s) = -isf_1(s) \in L_p(-1, 1)$ for a_n and $f(s) = f_2(s) \in L_p(-1, 1)$ for b_n . Hence it remains to show that $(-1)^n \sin \lambda_n = 1 + e_n(\tilde{g}_1)$ for some $\tilde{g}_1 \in L_p$ and that

$$(-1)^n \int_{-1}^1 f(s) e^{i\lambda_n s} ds = e_n(\tilde{g}_2)$$

for some $\tilde{g}_2 \in L_p$.

Since by Theorem 3.1 the eigenvalues λ_n satisfy the relation $\lambda_n = \pi(n + \frac{1}{2}) + \tilde{\lambda}_n$, where $\tilde{\lambda}_n = e_n(g_1)$ for some function $g_1 \in L_p$, we find that $\sin \lambda_n = (-1)^n \cos \tilde{\lambda}_n$, so that

$$(-1)^n \sin \lambda_n - 1 = \sum_{k=1}^{\infty} (-1)^k \frac{\tilde{\lambda}_n^{2k}}{(2k)!}.$$

Applying Proposition A.1 to the element $\mathbf{x} := (\tilde{\lambda}_n)_{n \in \mathbb{Z}} \in X_p$, we see that there exists a function $\tilde{g}_1$ in L_p such that the sum on the right-hand side of the above equality equals $e_n(\tilde{g}_1)$.

Changing the variables $s \mapsto 1 - 2t$ in the integral, we get

$$\begin{aligned} \int_{-1}^1 f(s) e^{i\lambda_n s} ds &= \int_0^1 2f(1 - 2t) e^{i\lambda_n(1-2t)} dt \\ &= (-1)^n \int_0^1 \tilde{f}(t) e^{i\tilde{\lambda}_n(1-2t)} e^{-2\pi nit} dt \end{aligned}$$

with $\tilde{f}(t) := 2if(1 - 2t)e^{-\pi it}$. Developing the function $e^{i\tilde{\lambda}_n(1-2t)}$ into the Taylor series and then changing summation and integration order (which is allowed in view of the absolute convergence of the Taylor series and the integral), we find that

$$(-1)^n \int_{-1}^1 f(s) e^{i\lambda_n s} ds = \sum_{k=0}^{\infty} \frac{\tilde{\lambda}_n^k}{k!} e_n(V^k \tilde{f}),$$

where V is the operator of multiplication by $i(1 - 2t)$. In virtue of Proposition A.1 and the fact that V has norm 1 in L_p , the right-hand side of the above equality gives the n -th Fourier coefficient of some function $\tilde{g}_2$ from L_p , and the proof is complete. $\square$

4. DERIVATION OF THE GLM AND THE KREIN EQUATION

Write $\mathbf{c}_n(\cdot) = \mathbf{c}(\cdot, \lambda_n)$ and $\alpha_n := \|\mathbf{c}_n\|_{\mathbb{H}}^{-2}$. Since the functions $\{\mathbf{c}_n\}_{n \in \mathbb{Z}}$ form an orthogonal basis of $\mathbb{H}$, we have

$$\text{s-lim}_{k \rightarrow \infty} \sum_{n=-k}^k \alpha_n \langle \cdot, \mathbf{c}_n \rangle \mathbf{c}_n = \mathcal{I},$$

where $\mathcal{I}$ is the identity operator in $\mathbb{H}$. On the other hand, $\mathbf{c}_n = (\mathcal{I} + \mathcal{K})\mathbf{v}_n$ with $\mathbf{v}_n(x) := \mathbf{c}_0(x, \lambda_n) = (\cos \lambda_n x, \sin \lambda_n x)^t$, so that the previous relation can be rewritten as

$$(\mathcal{I} + \mathcal{K}) \left[\text{s-lim}_{k \rightarrow \infty} \sum_{n=-k}^k \alpha_n \langle \cdot, \mathbf{v}_n \rangle \mathbf{v}_n \right] (\mathcal{I} + \mathcal{K})^* = \mathcal{I},$$

which implies

$$\text{s-lim}_{k \rightarrow \infty} \sum_{n=-k}^k \alpha_n \langle \cdot, \mathbf{v}_n \rangle \mathbf{v}_n = (\mathcal{I} + \mathcal{K})^{-1} (\mathcal{I} + \mathcal{K}^*)^{-1}.$$

Set

$$(4.1) \quad H(s) := \text{V.p.} \sum_{n=-\infty}^{\infty} \left(\alpha_n e^{-2\lambda_n s B} - e^{-\pi(2n+1)sB} \right),$$

where for $p = 1$ the summation is understood in the Cesàro sense, see Lemma 5.2. Observing that $\langle \cdot, \mathbf{c}_0(\cdot, \lambda) \rangle \mathbf{c}_0(\cdot, \lambda)$ is an integral operator with kernel

$$\frac{1}{2} [e^{-\lambda(x-t)B} + e^{-\lambda(x+t)B} J]$$

and that

$$\text{s-lim}_{k \rightarrow \infty} \sum_{n=-k}^k \langle \cdot, \mathbf{v}_{0,n} \rangle \mathbf{v}_{0,n} = \mathcal{I}$$

with $\mathbf{v}_{0,n} := \mathbf{c}_0(\cdot, \pi(n + \frac{1}{2}))$, we conclude that

$$(4.2) \quad (\mathcal{I} + \mathcal{K})^{-1} (\mathcal{I} + \mathcal{K}^*)^{-1} = \mathcal{I} + \mathcal{F},$$

where $\mathcal{F}$ is an integral operator in $\mathbb{H}$,

$$(\mathcal{F}\mathbf{u})(x) = \int_0^1 F(x, s) \mathbf{u}(s) ds,$$

with kernel

$$(4.3) \quad F(x, s) := \frac{1}{2} \left[H\left(\frac{x-t}{2}\right) + H\left(\frac{x+t}{2}\right) J \right].$$

Since $\mathcal{K}^*$ is an integral Volterra operator with upper-diagonal kernel, the operator $\mathcal{I} + \mathcal{K}^*$ is invertible and its inverse can be written in the form $\mathcal{I} + \tilde{\mathcal{K}}$, where $\tilde{\mathcal{K}}$ is an integral operator in $\mathbb{H}$ with upper-diagonal kernel. By (4.2) one gets

$$(\mathcal{I} + \mathcal{K})(\mathcal{I} + \mathcal{F}) = (\mathcal{I} + \mathcal{K}^*)^{-1} = \mathcal{I} + \tilde{\mathcal{K}};$$

spelling out this equality in terms of the kernels K and F for $x > t$, we arrive at the Gelfand–Levitan–Marchenko (GLM) equation

$$(4.4) \quad K(x, t) + F(x, t) + \int_0^x K(x, s) F(s, t) ds = 0.$$

If Q is continuous, then such is also K , and one has the formula [36, Lemma 12.1.1]

$$(4.5) \quad Q(x) = K(x, x)B - BK(x, x)$$

relating the potential and the kernel of the corresponding transformation operator. This suggests the following algorithm of solution of the inverse spectral problem: given the spectral data $\{(\lambda_n), (\alpha_n)\}$, one constructs first the kernel F via (4.3) and (4.1), then solves the GLM equation (4.4) for K , and, finally, recovers the potential via (4.5). However, if $Q \in L_p(\mathcal{M}_2)$, then relation (4.5) becomes meaningless since, by Theorem 2.3, in this case the kernel K belongs to $G_p(\mathcal{M}_2)$ but can be neither continuous nor well defined on subsets of $[0, 1] \times [0, 1]$ of Lebesgue measure zero.

It turns out that for $Q \in L_p(\mathcal{M}_2)$ the restriction $R(x, x)$ of the kernel R of (2.12) to the diagonal does determine a matrix-function with entries in $L_p(0, 1)$. If Q is continuous, then (4.5) together with (2.12) and the commutator relations

$$(4.6) \quad BR(x, t) = R(x, t)B, \quad BJ = -JB$$

yields the equality

$$(4.7) \quad Q(x) = R(x, x)JB.$$

Equation (4.7) retains sense also for $Q \in L_p(\mathcal{M}_2)$ as an equality in $L_p(\mathcal{M}_2)$ and thus can be used to recover Q in this situation.

Our next task is to explain how the kernel R can be determined from the spectral data. We do this by deriving below the Krein equation (4.9), an analogue of the GLM equation for R [10, 32].

Applying the operator B to equality (4.4) from both sides, we obtain its counterpart,

$$(4.8) \quad BK(x, t)B + BF(x, t)B + \int_0^x BK(x, s)F(s, t)B ds = 0.$$

With regard to the commutator relations (4.6) and $BH(x) = H(x)B$, we see that

$$\begin{aligned} BK(x, t)B &= \frac{1}{2}[-R(x, \frac{x-t}{2}) + R(x, \frac{x+t}{2})J], \\ BF(x, t)B &= \frac{1}{2}[-H(\frac{x-t}{2}) + H(\frac{x+t}{2})] \end{aligned}$$

and hence

$$\begin{aligned} K(x, t) + BK(x, t)B &= R(x, \frac{x+t}{2})J, \\ F(x, t) + BF(x, t)B &= H(\frac{x+t}{2})J. \end{aligned}$$

It also follows that

$$K(x, s)F(s, t) + BK(x, s)F(s, t)B = \frac{1}{2}[R(x, \frac{x-s}{2})H(\frac{s+t}{2})J + R(x, \frac{x+s}{2})JH(\frac{s-t}{2})].$$

Adding now (4.4) and (4.8), combining the above formulae in the resulting expression, and using the relation $JH(x) = H(-x)J$, we arrive at the equation

$$R(x, \frac{x+t}{2}) + H(\frac{x+t}{2}) + \int_0^x R(x, s)H(\frac{x+t}{2} - s) ds = 0, \quad 0 \leq t \leq x \leq 1.$$

Subtracting (4.8) from (4.4) and performing similar transformations, we arrive at the above formula with $\frac{x+t}{2}$ replaced by $\frac{x-t}{2}$, and both can now be combined together to give

$$R(x, t) + H(t) + \int_0^x R(x, s)H(t - s) ds = 0, \quad 0 \leq t \leq x \leq 1.$$

We see that the function $\tilde{R}(x, t) := R(x, x - t)$ satisfies the following Krein equation:

$$(4.9) \quad \tilde{R}(x, t) + H(x - t) + \int_0^x \tilde{R}(x, s)H(s - t) ds = 0.$$

We observe that as soon as a kernel H is given by (4.1) with λ_n and α_n obeying the proper asymptotics (guaranteeing that the series for H converges in $L_p(\mathcal{M}_2)$), the integral operator $\mathcal{H}$ with kernel $H(x - t)$ belongs to the algebra $\mathcal{G}_p(\mathcal{M}_2)$ introduced in Section 2 and $\mathcal{I} + \mathcal{H}$ is positive in $\mathbb{H}$ (see Lemma 5.4). The Krein equation (4.9) is then uniquely soluble and its solution belongs to $G_p(\mathcal{M}_2)$, see Appendix B. In particular, $\tilde{R}(x, 0) = R(x, x)$ is in $L_p(0, 1)$ entrywise indeed.

Remark 4.1. We notice that the GLM equation (4.4) is the even part of the Krein equation (4.9) in the sense that if $\tilde{R}$ is a solution to (4.9), then the function

$$(4.10) \quad K(x, t) := \frac{1}{2} [\tilde{R}(x, \frac{x-t}{2}) + \tilde{R}(x, \frac{x+t}{2})]$$

solves (4.4). Moreover, the condition $\mathcal{I} + \mathcal{H} > 0$ implies that the operator $\mathcal{I} + \mathcal{F}$ is positive in $\mathbb{H}$ and thus, in view of the results of Appendix B, guarantees that the GLM equation (4.4) with F of (4.3) is uniquely soluble for K and the solution belongs to $G_p(\mathcal{M}_2)$.

5. INVERSE SPECTRAL PROBLEM

The purpose of this section is, firstly, to show by limiting arguments that formula (4.7) remains valid if the matrix potential Q belongs to $L_p(\mathcal{M}_2)$ and, secondly, to justify the algorithm reconstructing the potential Q from the spectral data. Namely, we shall prove the following theorem, which constitutes the main result of the paper.

Theorem 5.1. *Assume that $\{(\lambda_n)_{n \in \mathbb{Z}}, (\mu_n)_{n \in \mathbb{Z}}\}$ is an arbitrary element of SD_p , $p \in [1, \infty)$. Then there exists a unique potential $Q \in L_p(\mathcal{M}_2)$ such that (λ_n) and (μ_n) are eigenvalues of the corresponding Dirac operators $\mathcal{A}_1$ and $\mathcal{A}_2$ respectively. The potential Q is equal to $R(x, x)JB = \tilde{R}(x, 0)JB$, where $\tilde{R}$ is the solution of the Krein equation (4.9) with H of (4.1) and α_n given by (3.7).*

The reconstruction algorithm proceeds as follows. Given an arbitrary element of SD_p , we construct functions ϕ and ψ via relations (3.6) and then determine the constants α_n by (3.7). Since the λ_n 's and μ_n 's interlace, it is easily seen that all α_n 's are positive. By virtue of the results of [27] there exist functions f_1 and f_2 in $L_p(-1, 1)$ such that

$$(5.1) \quad \begin{aligned} \phi(\lambda) &= \cos \lambda + \int_{-1}^1 f_1(s) e^{i\lambda s} ds, \\ \psi(\lambda) &= \sin \lambda + \int_{-1}^1 f_2(s) e^{i\lambda s} ds. \end{aligned}$$

Therefore the proof of Theorem 3.3 remains valid, and thus the numbers α_n satisfy the asymptotics $\alpha_n = 1 + e_n(g)$ for some $g \in L_p(0, 1)$. We shall now prove that the series (4.1) converges in $L_p(\mathcal{M}_2)$, so that the function H is well defined and belongs to $L_p(\mathcal{M}_2)$.

Lemma 5.2. *Assume that the numbers λ_n and α_n , $n \in \mathbb{Z}$, satisfy the asymptotics of Theorems 3.1 and 3.3 for some $p \in [1, \infty)$. Then the series (4.1) converges in the space $L_p((-1, 1), \mathcal{M}_2)$ (in the Cesàro sense if $p = 1$).*

Proof. Since the matrix B is skew-adjoint and has eigenvalues $\pm i$, it suffices to prove that the scalar series

$$(5.2) \quad h(s) := \text{V.p.} \sum_{n=-\infty}^{\infty} (\alpha_n e^{2\lambda_n i s} - e^{(2n+1)\pi i s})$$

converges in $L_p(-1, 1)$ (in the Cesàro sense if $p = 1$). We shall justify the convergence on $(0, 1)$; that on $(-1, 0)$ will then follow if we replace α_n with α_{-1-n} and λ_n with $-\lambda_{-1-n}$.

By assumption, $\alpha_n = 1 + e_n(g)$ for some $g \in L_p$, and classical theorems of harmonic analysis [28, Sec. I.2] show that the series

$$\text{V.p.} \sum_{n=-\infty}^{\infty} (\alpha_n - 1) e^{2n\pi i s}$$

converges in L_p to g in the required sense. It remains to establish that the series

$$(5.3) \quad \text{V.p.} \sum_{n=-\infty}^{\infty} \alpha_n (e^{2\lambda_n i s} - e^{(2n+1)\pi i s})$$

is convergent in L_p . We shall treat only the case $p = 1$; the arguments remain the same for $p > 1$ but with partial Cesàro sums replaced by the ordinary partial sums.

By the definition of summability in the sense of Cesàro we have to show that the sequence of partial sums

$$S_N(x) := \sum_{k=1}^N \sum_{n=-k}^k \alpha_n (e^{2\lambda_n i x} - e^{(2n+1)\pi i x})$$

is a Cauchy sequence in L_1 . Recalling that $\lambda_n = \pi(n + \frac{1}{2}) + \tilde{\lambda}_n$ with $\tilde{\lambda}_n = e_n(f)$ for some $f \in L_1$ and developing $e^{2\lambda_n i x}$ into Taylor series around $(2n+1)\pi i x$, we see that

$$\begin{aligned} S_N(x) &= \sum_{k=1}^N \sum_{n=-k}^k \alpha_n e^{(2n+1)\pi i x} \sum_{m=1}^{\infty} \frac{(2\tilde{\lambda}_n i x)^m}{m!} \\ &= e^{\pi i x} \sum_{m=1}^{\infty} \frac{(2i x)^m}{m!} \sum_{k=1}^N \sum_{n=-k}^k \alpha_n \tilde{\lambda}_n^m e^{2n\pi i x} \\ &= e^{\pi i x} \sum_{m=1}^{\infty} \frac{(2i x)^m}{m!} \sigma_N(x, f_m), \end{aligned}$$

where

$$\sigma_N(x, f_m) := \sum_{k=1}^N \sum_{n=-k}^k e_n(f_m) e^{2n\pi i x}$$

is the partial Cesàro sum for the function

$$f_m := \underbrace{f * \cdots * f}_m + \underbrace{f * \cdots * f}_m * g, \quad m \geq 1$$

(see Appendix A). As is well known (cf. [28, Ch. II]), for any $f \in L_1$, the partial Cesàro sums $\sigma_N(\cdot, f)$ converge to f in L_1 and $\|\sigma_N(\cdot, f)\|_{L_1} \leq \|f\|_{L_1}$. Since

$$\|f_m\|_{L_1} \leq (1 + \|g\|_{L_1}) \|f\|_{L_1}^m,$$

the Lebesgue dominated convergence theorem shows that the limit

$$\lim_{N \rightarrow \infty} S_N(x) = e^{-\pi i x} \sum_{m=1}^{\infty} \frac{(2ix)^m}{m!} f_m(x)$$

exists in L_1 , whence the series (5.3) converges in L_1 in the Cesàro sense. The lemma is proved. $\square$

Next we show that the GLM equation (4.4) and the Krein equation (4.9) have unique solutions that belong to the space $G_p(\mathcal{M}_2)$. To this end it suffices to show (see Appendix B and Remark 4.1 for details) that the operator $\mathcal{I} + \mathcal{H}$ is positive in $\mathbb{H}$, where $\mathcal{H}$ is a Wiener–Hopf operator given by

$$\mathcal{H}\mathbf{u}(x) := \int_0^1 H(x-s)\mathbf{u}(s) ds.$$

As a preliminary, we show that certain systems of functions form Riesz bases in L_2 or $\mathbb{H}$.

Lemma 5.3. *Assume that the sequence (λ_n) is as in Theorem 5.1. Then*

- (a) *the system $(e^{i\lambda_n x})_{n \in \mathbb{Z}}$ forms a Riesz basis of $L_2(-1, 1)$;*
- (b) *the system $(\mathbf{v}_n)_{n \in \mathbb{Z}}$, with $\mathbf{v}_n := \mathbf{c}_0(\cdot, \lambda_n)$, forms a Riesz basis of $\mathbb{H}$.*

Proof. In view of relation (3.3) the numbers λ_k are zeros of the exponential function of sine type with indicator diagram $[-1, 1]$ [34], so that item (a) follows from [6, Proposition II.4.3].

Assume now that $\mathbf{g} := (g_1, g_2)^t$ is an arbitrary element of $\mathbb{H}$. We extend g_1 and g_2 to functions $\tilde{g}_1$ and $\tilde{g}_2$ on $(-1, 1)$ in the even and odd way, respectively, and set $\tilde{g} := \tilde{g}_1 + i\tilde{g}_2$. By (a), there exists a unique sequence $(a_n)_{n \in \mathbb{Z}}$ in ℓ_2 , for which

$$\tilde{g} = \sum_{n=-\infty}^{\infty} a_n e^{i\lambda_n x},$$

the series being convergent in $L_2(-1, 1)$. Taking the even and odd parts of the above equality, we arrive at the relations

$$\tilde{g}_1(x) = \sum_{n=-\infty}^{\infty} a_n \cos(\lambda_n x), \quad i\tilde{g}_2(x) = i \sum_{n=-\infty}^{\infty} a_n \sin(\lambda_n x),$$

i.e., at the representation

$$\mathbf{g} = \sum_{n=-\infty}^{\infty} a_n \mathbf{v}_n$$

in $\mathbb{H}$. We observe that there is a constant C independent of $\tilde{g}$ such that

$$C^{-1} \sum_{n=-\infty}^{\infty} |a_n|^2 \leq \|\tilde{g}\|_{L_2}^2 = 2\|\mathbf{g}\|_{\mathbb{H}}^2 \leq C \sum_{n=-\infty}^{\infty} |a_n|^2.$$

Hence the system $(\mathbf{v}_n)_{n \in \mathbb{Z}}$ is a Riesz basis of $\mathbb{H}$. $\square$

Lemma 5.4. *Assume that $\{(\lambda_n), (\mu_n)\}$ is an arbitrary element of SD_p and that the function H is constructed as explained above. Then the corresponding operator $\mathcal{I} + \mathcal{H}$ is positive in $\mathbb{H}$.*

Proof. We notice that the matrix B is skew-adjoint and has eigenvalues $\pm i$. It is clear that the subspaces $\ker(B \pm i) \otimes L_2(0, 1)$ are invariant subspaces of $\mathcal{H}$. Therefore $\mathcal{H}$ is unitarily equivalent to the direct sum $\mathcal{H}_+ \oplus \mathcal{H}_-$, where

$$(\mathcal{H}_\pm g)(s) = \int_0^1 h(\pm(x-s))g(s) ds, \quad g \in L_2,$$

and h is the function of (5.2). Thus positivity of $\mathcal{I} + \mathcal{H}$ is equivalent to that of both $I + \mathcal{H}_+$ and $I + \mathcal{H}_-$. We shall prove only that the operator $I + \mathcal{H}_+$ is positive in L_2 ; the positivity of the other one is established analogously.

Since the systems $(e^{(2n+1)\pi is})_{n \in \mathbb{Z}}$ and $(e^{2\lambda_n is})_{n \in \mathbb{Z}}$ constitute respectively an orthonormal and a Riesz basis of L_2 (cf. Lemma 5.3 and [6, Sect. II.4.2]), it is easy to see that

$$\begin{aligned} ((I + \mathcal{H}_+)f, f) &= (f, f) + \lim_{k \rightarrow \infty} \sum_{n=-k}^k [\alpha_n |(f, e^{2\lambda_n is})|^2 - |(f, e^{(2n+1)\pi is})|^2] \\ &= \sum_{n=-\infty}^{\infty} \alpha_n |(f, e^{2\lambda_n is})|^2. \end{aligned}$$

Since the numbers α_n are uniformly bounded away from zero, we conclude that there is a $C > 0$ such that $((I + \mathcal{H}_+)f, f) \geq C\|f\|^2$ for all $f \in L_2$, i.e., the operator $I + \mathcal{H}_+$ is (uniformly) positive in L_2 . $\square$

According to Theorem B.2, positivity of the operator $\mathcal{I} + \mathcal{H}$ implies that the GLM equation (4.4) with F given by (4.3) and the Krein equation (4.9) have unique solutions $K, \tilde{R} \in G_p(\mathcal{M}_2)$. We have shown in Section 4 that in the smooth case the solution K is the kernel of the transformation operator $\mathcal{I} + \mathcal{K}$ for the pair $(\tilde{\mathcal{A}}_Q, \tilde{\mathcal{A}}_0)$ with $Q := \tilde{R}(\cdot, 0)JB$. Based on this result, we treat here the general case by a limiting procedure.

Theorem 5.5. *Assume that $\{(\lambda_n), (\mu_n)\}$ is an arbitrary element of SD_p and that $H \in L_p(\mathcal{M}_2)$ is a function of (4.1) constructed as explained above. Let also K and $\tilde{R}$ be the solutions of the GLM equation (4.4) and the Krein equation (4.9) respectively. Denote by $\mathcal{K}$ the integral operator with kernel K . Then there exist a unique $Q \in L_p(\mathcal{M}_2)$ —namely, $Q = \tilde{R}(\cdot, 0)JB$ —such that the operator $\mathcal{I} + \mathcal{K}$ is the transformation operator for the pair $\tilde{\mathcal{A}}_Q$ and $\tilde{\mathcal{A}}_0$.*

Proof. We shall approximate the function H in the norm of $L_p(\mathcal{M}_2)$ by a sequence $(H_l)_{l=1}^\infty$ of real-valued, smooth (say, infinitely differentiable) $\mathcal{M}_2$ -valued functions so that the following holds:

- (a) for every $l \in \mathbb{N}$, the GLM equation (4.4) with H replaced by H_l has a unique solution K_l , and the corresponding integral operators $\mathcal{K}_l$ converge to $\mathcal{K}$ as $l \rightarrow \infty$ in the uniform operator topology of $\mathbb{H}$;
- (b) for every $l \in \mathbb{N}$, there exists $Q_l \in L_p(\mathcal{M}_2)$ of the form (1.1) such that $\mathcal{I} + \mathcal{K}_l$ is a transformation operator for the pair $\tilde{\mathcal{A}}_{Q_l}$ and $\tilde{\mathcal{A}}_0$;
- (c) the matrix-functions Q_l converge to $Q := \tilde{R}(\cdot, 0)JB$ in $L_p(\mathcal{M}_2)$.

If (a)–(c) hold, then by Theorems 2.3 and 2.4 the operators $\mathcal{I} + \mathcal{K}_l$ converge in $\mathcal{G}_p(\mathcal{M}_2)$ (and hence in the uniform operator topology of $\mathbb{H}$) to an operator $\mathcal{I} + \mathcal{K}_Q$, which is the transformation operator for the pair $(\tilde{\mathcal{A}}_Q, \tilde{\mathcal{A}}_0)$. Thus $\mathcal{K} = \mathcal{K}_Q$ yielding the result. The uniqueness of Q is obvious.

The details are as follows. Using (λ_n) and (μ_n) , we construct the sequence of constants α_n and set

$$H_l(s) := \sum_{n=-l}^l \left(\alpha_n e^{-2\lambda_n s B} - e^{-\pi(2n+1)sB} \right)$$

(for $p = 1$, we replace H_l by the corresponding partial Cesàro sum $\frac{1}{l} \sum_{k=1}^l H_k$); then $H_l \rightarrow H$ in $L_p(\mathcal{M}_2)$ as $l \rightarrow \infty$ by Lemma 5.2.

We observe that this choice of H_l corresponds to setting $\alpha_n = 1$ and $\lambda_n = \pi(n + \frac{1}{2})$ for all n with $|n| > l$, so that in view of Lemma 5.4 the Wiener–Hopf operators $\mathcal{H}_l$ with symbol H_l satisfy the condition $\mathcal{J} + \mathcal{H}_l > 0$. Hence by Corollary B.3 and Remark 4.1 the GLM equation (4.4) with H replaced by H_l has a unique solution K_l . This solution K_l belongs to $G_p(\mathcal{M}_2)$, and hence the corresponding integral operator $\mathcal{K}_l$ is bounded. Since the relation $H_l \rightarrow H$ in $L_p(\mathcal{M}_2)$ as $l \rightarrow \infty$ implies that $\mathcal{H}_l \rightarrow \mathcal{H}$ in $\mathcal{G}_p(\mathcal{M}_2)$, we conclude that $\mathcal{K}_l \rightarrow \mathcal{K}$ as $l \rightarrow \infty$ in the topology of the space $\mathcal{G}_p(\mathcal{M}_2)$ —and thus in the uniform operator topology of $\mathbb{H}$, see Appendix B. This establishes (a).

Moreover, by the well-known result for continuous potentials [36, Ch. 12.3–4] the operator $\mathcal{J} + \mathcal{K}_l$ is the transformation operator for the pair $(\tilde{\mathcal{A}}_{Q_l}, \tilde{\mathcal{A}}_0)$ with $Q_l(x) := K_l(x, x)B - BK_l(x, x)$. As was explained in Section 4, this yields the relation $Q_l(\cdot) = \tilde{R}_l(\cdot, 0)JB$, where $\tilde{R}_l$ is the solution to the Krein equation (4.9) for $H = H_l$. Thus (b) is fulfilled.

To prove (c), we observe that, according to the results of Appendix B, the solution $\tilde{R}$ to the Krein equation (4.9) depends continuously in the norm of the space $G_p(\mathcal{M}_2)$ on the function $H \in L_p(\mathcal{M}_2)$. Therefore the sequence $\tilde{R}_l(\cdot, 0)$ converges in $L_p(\mathcal{M}_2)$ to the function $\tilde{R}(\cdot, 0)$. The proof is complete. $\square$

The last step of the reconstruction procedure is to show that the numbers λ_n and μ_n we have started with are the very eigenvalues of the operators $\mathcal{A}_1$ and $\mathcal{A}_2$ with the potential Q just found.

Since the solution K to the GLM equation (4.4) generates a transformation operator $\mathcal{J} + \mathcal{K}$ for the pair $(\tilde{\mathcal{A}}_Q, \tilde{\mathcal{A}}_0)$, the functions $\mathbf{c}(\cdot, \lambda) := (\mathcal{J} + \mathcal{K})\mathbf{c}_0(\cdot, \lambda)$ belong to $\text{dom } \tilde{\mathcal{A}}_Q$ and satisfy the relation $\tilde{\mathcal{A}}_Q \mathbf{c}(\cdot, \lambda) = \lambda \mathbf{c}(\cdot, \lambda)$. We set $\mathbf{c}_k := \mathbf{c}(\cdot, \lambda_k)$ and show that these functions are orthogonal and that the α_k are the corresponding norming constants.

Lemma 5.6. *The system of functions $\{\mathbf{c}_k\}_{k \in \mathbb{Z}}$ is an orthogonal basis of $\mathbb{H}$. Moreover, for the above numbers α_k (defined at the beginning of this section), we have*

$$\langle \mathbf{c}_k, \mathbf{c}_l \rangle = \alpha_k^{-1} \delta_{kl},$$

where δ_{kl} is the Kronecker delta.

Proof. Denoting by $\mathbf{v}_n$ the function $\mathbf{c}_0(\cdot, \lambda_n)$ and recalling that the integral operator $\mathcal{F}$ with kernel F of (4.3) is related to $\mathcal{K}$ by $(\mathcal{J} + \mathcal{K})(\mathcal{J} + \mathcal{F})(\mathcal{J} + \mathcal{K})^* = \mathcal{J}$ (see Appendix B for details), we conclude that

$$\langle \mathbf{c}_k, \mathbf{c}_l \rangle = \langle (\mathcal{J} + \mathcal{K})^*(\mathcal{J} + \mathcal{K})\mathbf{v}_k, \mathbf{v}_l \rangle = \langle (\mathcal{J} + \mathcal{F})^{-1}\mathbf{v}_k, \mathbf{v}_l \rangle.$$

Reverting the arguments of Section 4, we see that the operator $\mathcal{J} + \mathcal{F}$ is equal to

$$\text{s-lim}_{k \rightarrow \infty} \sum_{n=-k}^k \alpha_n \langle \cdot, \mathbf{v}_n \rangle \mathbf{v}_n.$$

Since the sequence $(\mathbf{v}_n)_{n \in \mathbb{Z}}$ is a Riesz basis of $\mathbb{H}$ in view of Lemma 5.3, the inverse of $\mathcal{J} + \mathcal{F}$ can be represented as

$$(\mathcal{J} + \mathcal{F})^{-1} = \text{s-lim}_{n \rightarrow \infty} \sum_{m=-n}^n \alpha_m^{-1} \langle \cdot, \tilde{\mathbf{v}}_m \rangle \tilde{\mathbf{v}}_m,$$

where $(\tilde{\mathbf{v}}_m)$ is a basis biorthogonal to $(\mathbf{v}_m)$ (see [19, Ch. VI]). Therefore

$$\langle (\mathcal{J} + \mathcal{F})^{-1} \mathbf{v}_k, \mathbf{v}_l \rangle = \text{s-lim}_{n \rightarrow \infty} \sum_{m=-n}^n \alpha_m^{-1} \langle \mathbf{v}_k, \tilde{\mathbf{v}}_m \rangle \langle \tilde{\mathbf{v}}_m, \mathbf{v}_l \rangle = \sum_{m=-\infty}^{\infty} \alpha_m^{-1} \delta_{km} \delta_{ml} = \alpha_k^{-1} \delta_{kl}.$$

Completeness of the system $\{\mathbf{c}_n\}_{n \in \mathbb{Z}}$ immediately follows from the fact that the system $\{\mathbf{v}_n\}_{n \in \mathbb{Z}}$ is complete and that $\mathcal{J} + \mathcal{K}$ is a homeomorphism of $\mathbb{H}$, and the lemma is proved. $\square$

Next we show that the numbers λ_n are indeed the eigenvalues of the operator $\mathcal{A}_1$. According to what was said above, it suffices to show that $c_1(1, \lambda_k) = 0$. For the operator $\tilde{\mathcal{A}}_Q$, one has the Green formula

$$\langle \tilde{\mathcal{A}}_Q \mathbf{c}(\cdot, \lambda), \mathbf{c}(\cdot, \mu) \rangle - \langle (\mathbf{c}(\cdot, \lambda), \tilde{\mathcal{A}}_Q \mathbf{c}(\cdot, \mu)) \rangle = c_2(1, \lambda) c_1(1, \mu) - c_1(1, \lambda) c_2(1, \mu);$$

taking therein $\lambda = \lambda_k$ and $\mu = \lambda_n$ and using the previous lemma, we arrive at the relation

$$(5.4) \quad c_2(1, \lambda_k) c_1(1, \lambda_n) = c_1(1, \lambda_k) c_2(1, \lambda_n).$$

Assume first that none of the numbers $c_1(1, \lambda_k)$ vanishes. Relation (5.4) implies that there is a constant γ such that, for all $k \in \mathbb{Z}$,

$$c_2(1, \lambda_k) / c_1(1, \lambda_k) = \gamma.$$

Then $\mathbf{c}_k$ are eigenvectors corresponding to the eigenvalues λ_k of the operator that is the restriction of $\tilde{\mathcal{A}}_Q$ by the boundary condition $u_2(1) = \gamma u_1(1)$. In other words, the numbers λ_n are zeros of the function

$$c_2(1, \lambda) - \gamma c_1(1, \lambda) = \sin \lambda - \gamma \cos \lambda + \int_{-1}^1 \tilde{f}(s) e^{i\lambda s} ds$$

for some $\tilde{f} \in L_p(-1, 1)$. However, the standard arguments based on Rouché's theorem (see, e.g., [41, Ch. 1.3]) show that the zeros $\tilde{\lambda}_n$ of the function $c_2(1, \lambda) - \gamma c_1(1, \lambda)$ obey the different asymptotics $\tilde{\lambda}_n = \pi n + \arctan \gamma + o(1)$, which leads to a contradiction.

Therefore there is a $k \in \mathbb{Z}$ such that $c_1(1, \lambda_k) = 0$. Then $c_2(1, \lambda_k) \neq 0$ due to the uniqueness of solutions to the equation $\ell_Q(\mathbf{u}) = \lambda_k \mathbf{u}$, and relation (5.4) implies that $c_2(1, \lambda_n) = 0$ for all $n \in \mathbb{Z}$. In other words, the functions $\mathbf{c}_n$ are the eigenvectors of the operator $\mathcal{A}_1$ corresponding to the eigenvalues λ_n . Since by Lemma 5.6 the system $\{\mathbf{c}_n\}_{n \in \mathbb{Z}}$ is complete in $\mathbb{H}$, the operator $\mathcal{A}_1$ has no other eigenvalues.

It remains to prove that μ_n are the eigenvalues of $\mathcal{A}_2$. We denote by $\tilde{\mu}_n$ these eigenvalues and construct the corresponding function $\tilde{\psi}$ of (3.6). Recalling expression (3.7), we conclude that $\psi(\lambda_k) = \tilde{\psi}(\lambda_k)$ for all $k \in \mathbb{Z}$. Since the function $\tilde{\psi}$ has the representation

$$\tilde{\psi}(\lambda) = \sin \lambda + \int_{-1}^1 \tilde{f}_2(s) e^{i\lambda s} ds$$

for some $\tilde{f}_2 \in L_p(-1, 1)$ (see the proof of Theorem 3.1) and ψ has a similar representation with some $f_2 \in L_p(-1, 1)$ instead of $\tilde{f}$ by (5.1), we see that the function $\omega := f_2 - \tilde{f}_2$ is such that

$$\int_{-1}^1 \omega(s) e^{i\lambda_n s} ds = 0$$

for all $n \in \mathbb{Z}$. Recalling that the system of functions $\{e^{i\lambda_n s}\}_{n \in \mathbb{Z}}$ is closed in $L_p(-1, 1)$ (this follows from [35, Theorem III] for $p > 1$ and from [27, Lemma 3.3] for $p = 1$), we conclude that $\omega = 0$. Thus the numbers $\mu_n = \tilde{\mu}_n$ are the eigenvalues of the operator $\mathcal{A}_2$, and the reconstruction procedure is complete. As the spectral data determine the function H (and thus the transformation operator $\mathcal{J} + \mathcal{K}$) unambiguously, the potential Q is unique. The proof of Theorem 5.1 is complete.

We observe that, in passing, we have solved the inverse spectral problem of reconstructing the potential Q of the Dirac operator from the spectrum of $\mathcal{A}_1$ and the corresponding sequence of norming constants. Namely, the following is true.

Theorem 5.7. *Sequences of real numbers $(\lambda_n)_{n \in \mathbb{Z}}$ and positive numbers $(\alpha_n)_{n \in \mathbb{Z}}$ are respectively the sequences of eigenvalues and norming constants of an operator $\mathcal{A}_1$ for some $Q \in L_p(\mathcal{M}_2)$, $p \in [1, \infty)$, if and only if the following holds:*

- (i) *the numbers λ_n strictly increase and obey the asymptotics of Theorem 3.1;*
- (ii) *the numbers $\alpha_n > 0$ obey the asymptotics of Theorem 3.3.*

If (i) and (ii) hold, then Q is given by $R(x, x)JB = \tilde{R}(x, 0)JB$, where $\tilde{R}$ is the solution of the Krein equation (4.9) with H of (4.1).

The reconstruction algorithm proceeds as the previous one, except that the first and the last step (related to the spectrum (μ_n)) should be omitted.

In a similar manner we can also treat the inverse spectral problem for the operator $\mathcal{A}_2$ (or operators corresponding to arbitrary separated boundary conditions).

Acknowledgements. The authors express their gratitude to DFG for financial support of the project 436 UKR 113/79. The second author gratefully acknowledges the financial support of the Alexander von Humboldt Foundation. The second and the third authors thank the Institute for Applied Mathematics of Bonn University for the warm hospitality.

APPENDIX A. FOURIER TRANSFORM IN $L_p(0, 1)$

For any $f \in L_p(0, 1)$, we denote by $e_n(f)$, $n \in \mathbb{Z}$, its n -th Fourier coefficients, i.e.,

$$e_n(f) = \int_0^1 f(x) e^{-2\pi n i x} dx.$$

We also denote by $\mathbf{e}(f)$ the sequence $(e_n(f))_{n \in \mathbb{Z}}$ and put

$$X_p := \{\mathbf{e}(f) \mid f \in L_p(0, 1)\}.$$

The vector space X_p is algebraically embedded into $\ell_\infty(\mathbb{Z})$ and becomes a Banach space under the induced norm

$$\|\mathbf{e}(f)\|_{X_p} := \|f\|_{L_p}.$$

For any $\mathbf{x} = (x_n)$ and $\mathbf{y} = (y_n)$ in $\ell_\infty(\mathbb{Z})$ we shall denote by $\mathbf{xy}$ the entrywise product of $\mathbf{x}$ and $\mathbf{y}$, i.e., the element of $\ell_\infty(\mathbb{Z})$ with the n -th entry $x_n y_n$.

Proposition A.1. X_p is a commutative Banach algebra under the entrywise multiplication, i.e.,

$$(A.1) \quad \|\mathbf{xy}\|_{X_p} \leq \|\mathbf{x}\|_{X_p} \|\mathbf{y}\|_{X_p}.$$

Indeed, inequality (A.1) follows from the fact that $e_n(f)e_n(g) = e_n(f * g)$, where

$$(f * g)(x) := \int_0^1 f(x-t)g(t) dt$$

is the convolution of f and g (f being periodically extended to $(-1, 1)$ by $f(x+1) = f(x)$, $x \in (0, 1)$), and from the inequality

$$\|f * g\|_{L_p} \leq \|f\|_{L_p} \|g\|_{L_p}.$$

The following statement is an analogue of the well-known Wiener lemma.

Proposition A.2. Assume that $f \in L_p(0, 1)$, where $p \in [1, \infty)$. If $1 + e_n(f) \neq 0$ for all $n \in \mathbb{N}$, then there exists a function $g \in L_p(0, 1)$ such that

$$(1 + e_n(f))^{-1} = 1 + e_n(g).$$

Proof. To begin with, we adjoin to X_p the unit element δ (with all components equal to 1) and denote the resulting unital algebra by $\widehat{X}_p$. Assume that the assumptions of the lemma hold and denote by x an element of $\widehat{X}_p$ with components $x_n := 1 + e_n(f)$. We shall prove below that x is invertible in $\widehat{X}_p$; since $e_n(f) \rightarrow 0$ as $n \rightarrow \infty$, this will imply that $x^{-1} = \delta + y$ for some $y \in X_p$ as required.

As is well known [46], the element x is invertible in the unital Banach algebra $\widehat{X}_p$ if and only if x does not belong to any maximal ideal of $\widehat{X}_p$. Proceeding by contradiction, assume that there exists a maximal ideal $\mathfrak{m}$ of $\widehat{X}_p$ containing x . Since $\widehat{X}_p$ includes all finite sequences and none of x_n vanishes, $\mathfrak{m}$ also contains all finite sequences. Finite sequences form a dense subset of X_p because the set of all trigonometric polynomials is dense in $L_p(0, 1)$. Recalling that maximal ideals are closed, we conclude that $X_p \subset \mathfrak{m}$. Next we observe that X_p is a proper subset of $\mathfrak{m}$ (e.g., x belongs to $\mathfrak{m} \setminus X_p$) and that X_p has codimension 1 in $\widehat{X}_p$. Hence $\mathfrak{m} = \widehat{X}_p$, which contradicts our assumption that $\mathfrak{m}$ is a maximal ideal of $\widehat{X}_p$. As a result, x is not contained in any maximal ideal of $\widehat{X}_p$ and thus is invertible in $\widehat{X}_p$ indeed. The lemma is proved. $\square$

APPENDIX B. THE GLM EQUATION AND FACTORISATION OF FREDHOLM OPERATORS

In this appendix, we shall explain relationships between solubility of the GLM equation and factorisation of related Fredholm operators in some special algebras. We refer the reader to the books [18, 20] for related concepts and basic facts.

Write $\mathbb{H} := L_2((0, 1), \mathbb{C}^2)$ and denote by $\mathcal{B}$ (by $\mathcal{B}_\infty$) the Banach algebra of all bounded (compact) operators in $\mathbb{H}$. Denote also by P_t , $t \in [0, 1]$, the operator in $\mathbb{H}$ of multiplication by $\chi_{[0, t]}$, the characteristic function of the interval $[0, t]$. Set

$$\begin{aligned} \mathcal{B}_\infty^+ &:= \{B \in \mathcal{B}_\infty \mid \forall t \in [0, 1], \quad P_t B (I - P_t) = 0\}, \\ \mathcal{B}_\infty^- &:= \{B \in \mathcal{B}_\infty \mid \forall t \in [0, 1], \quad (I - P_t) B P_t = 0\}; \end{aligned}$$

then $\mathcal{B}_\infty^\pm$ are closed subspaces of $\mathcal{B}_\infty$ and $\mathcal{B}_\infty^+ \cap \mathcal{B}_\infty^- = \{0\}$. We also observe that the operators in $\mathcal{B}_\infty^\pm$ are Volterra ones.

Recall that $\mathcal{G}_p(\mathcal{M}_2)$, $p \in [1, \infty)$, stands for the algebra in $\mathcal{B}_\infty$ of integral operators over $(0, 1)$ with kernels in the space $G_p(\mathcal{M}_2)$ introduced in Section 2. The sets

$$\mathcal{G}_p^+(\mathcal{M}_2) = \mathcal{G}_p(\mathcal{M}_2) \cap \mathcal{B}_\infty^+, \quad \mathcal{G}_p^-(\mathcal{M}_2) := \mathcal{G}_p(\mathcal{M}_2) \cap \mathcal{B}_\infty^-,$$

are subalgebras of $\mathcal{G}_p(\mathcal{M}_2)$ consisting of operators with lower- and upper-triangular kernels respectively and $\mathcal{G}_p(\mathcal{M}_2) = \mathcal{G}_p^+(\mathcal{M}_2) \dot{+} \mathcal{G}_p^-(\mathcal{M}_2)$.

We say that an operator $\mathcal{I} + \mathcal{L}$, $\mathcal{L} \in \mathcal{G}_p(\mathcal{M}_2)$, admits a factorization in $\mathcal{G}_p(\mathcal{M}_2)$ if

$$(B.1) \quad \mathcal{I} + \mathcal{L} = (\mathcal{I} + \mathcal{K}^+)^{-1}(\mathcal{I} + \mathcal{K}^-)^{-1}$$

with some $\mathcal{K}^\pm \in \mathcal{G}_p^\pm(\mathcal{M}_2)$.

The following two theorems were established in [43, 44] for the space $L_2(0, 1)$, however, their generalisation to our situation is straightforward.

Theorem B.1. *If $\mathcal{I} + \mathcal{L}$ admits a factorization in $\mathcal{G}_p(\mathcal{M}_2)$, then the operators $\mathcal{K}^\pm = \mathcal{K}^\pm(\mathcal{L})$ are unique. Moreover, the set Φ_p of operators $\mathcal{L} \in \mathcal{G}_p(\mathcal{M}_2)$, for which the operator $\mathcal{I} + \mathcal{L}$ is factorisable, is open in $\mathcal{G}_p(\mathcal{M}_2)$, and the functions*

$$\Phi_p \ni \mathcal{L} \mapsto \mathcal{K}^\pm(\mathcal{L}) \in \mathcal{G}_p(\mathcal{M}_2)$$

are continuous.

Theorem B.2. *Assume that $\mathcal{L} \in \mathcal{G}_p(\mathcal{M}_2)$. For the operator $\mathcal{I} + \mathcal{L}$ to admit a factorisation in $\mathcal{G}_p(\mathcal{M}_2)$, it is necessary and sufficient that the operators $I + P_t \mathcal{L} P_t$ have a trivial kernel in $\mathbb{H}$ for each $t \in [0, 1]$.*

We remark that for a self-adjoint operator $\mathcal{L} \in \mathcal{G}_p(\mathcal{M}_2)$ the requirement that the operators $I + P_t \mathcal{L} P_t$ have a trivial kernel in $\mathbb{H}$ for all $t \in [0, 1]$ is equivalent to positivity of $\mathcal{I} + \mathcal{L}$ in $\mathbb{H}$.

Assume that $\mathcal{I} + \mathcal{L}$ is factorisable in $\mathcal{G}_p(\mathcal{M}_2)$, so that (B.1) holds. Applying $\mathcal{I} + \mathcal{K}^+$ to both sides of this equality and using the fact that $(\mathcal{I} + \mathcal{K}^-)^{-1} = \mathcal{I} + \tilde{\mathcal{K}}^-$ for some $\tilde{\mathcal{K}}^- \in \mathcal{B}_\infty^-$, we derive the relation

$$(B.2) \quad \mathcal{K}^+ + \mathcal{P}^+ \mathcal{L} + \mathcal{P}^+(\mathcal{K}^+ \mathcal{L}) = 0,$$

where $\mathcal{P}^+$ denotes the projection operator of $\mathcal{G}_p(\mathcal{M}_2)$ onto $\mathcal{G}_p^+(\mathcal{M}_2)$ parallel to $\mathcal{G}_p^-(\mathcal{M}_2)$. This relation is an abstract analogue of the Gelfand–Levitan–Marchenko (GLM) equation; indeed, in terms of the kernels K^+ and L of the operators $\mathcal{K}^+$ and $\mathcal{L}$ we get

$$(B.3) \quad K^+(x, t) + L(x, t) + \int_0^x K^+(x, s) L(s, t) ds = 0, \quad 0 \leq t \leq x \leq 1,$$

cf. (4.4) and (4.9).

We see that if an operator $\mathcal{I} + \mathcal{L}$ is factorisable in $\mathcal{G}_p(\mathcal{M}_2)$, then the abstract GLM equation (B.2) has a solution $\mathcal{K}^+ = \mathcal{K}^+(\mathcal{L}) \in \mathcal{G}_p^+(\mathcal{M}_2)$. Conversely, if $\mathcal{K}^+ \in \mathcal{G}_p^+(\mathcal{M}_2)$ is a solution of equation (B.2), then $\mathcal{P}^+(\mathcal{K}^+ + \mathcal{L} + \mathcal{K}^+ \mathcal{L}) = 0$, i.e., $\mathcal{K} := \mathcal{K}^+ + \mathcal{L} + \mathcal{K}^+ \mathcal{L}$ belongs to $\mathcal{G}_p^-(\mathcal{M}_2)$, and the relation

$$(\mathcal{I} + \mathcal{K}^+)(\mathcal{I} + \mathcal{L}) = \mathcal{I} + \mathcal{K}$$

holds. Since $\mathcal{I} + \mathcal{K}$ has the form $(\mathcal{I} + \mathcal{K}^-)^{-1}$ for $\mathcal{K}^- := (\mathcal{I} + \mathcal{K})^{-1} - \mathcal{I} \in \mathcal{G}_p^-(\mathcal{M}_2)$, we conclude that $\mathcal{I} + \mathcal{L}$ is factorisable in $\mathcal{G}_p(\mathcal{M}_2)$.

Summing up, we derive the following assertion on the solubility of the GLM equation (B.3).

Corollary B.3. Assume that $\mathcal{L}$ is a selfadjoint operator in $\mathcal{G}_p(\mathcal{M}_2)$ with kernel L such that $\mathcal{J} + \mathcal{L}$ is positive. Then equation (B.3) is uniquely soluble, and the solution K^+ belongs to $G_p(\mathcal{M}_2)$ and depends continuously therein on $L \in G_p(\mathcal{M}_2)$.

REFERENCES

- [1] M. J. Ablowitz, D. J. Kaup, A. C. Newell, and H. Segur, Nonlinear-evolution equations of physical significance, *Phys. Rev. Lett.* **31** (1973), 125–127.
- [2] M. J. Ablowitz, D. J. Kaup, A. C. Newell, and H. Segur, The inverse scattering transform-Fourier analysis for nonlinear problems, *Studies in Appl. Math.* **53** (1974), no. 4, 249–315.
- [3] S. Albeverio, R. Hryniv, and Ya. Mykytyuk, Inverse spectral problems for Sturm–Liouville operators in impedance form, *J. Funct. Anal.* **222** (2005), 143–177.
- [4] S. Albeverio, L. Nizhnik, and V. Tarasov, Inverse scattering problem for a Dirac system with nonstationary point interactions, *Inverse Problems* **20** (2004), no. 3, 799–813.
- [5] T. N. Arutyunyan, Asymptotics of the Weyl–Titchmarsh function and the inverse problem for the Dirac system, *Izv. Akad. Nauk Armyan. SSR Ser. Mat.* **24** (1989), no. 4, 327–336, 416 (Russian); Engl. transl.: *Soviet J. Contemporary Math. Anal.* **24** (1989), no. 4, 15–24.
- [6] S. A. Avdonin and S. A. Ivanov, *Families of Exponentials: the Method of Moments in Controllability Problems for Distributed Parameter Systems*, Cambridge Univ. Press, Cambridge, 1995.
- [7] S. Clark and F. Gesztesy, Weyl–Titchmarsh M -function asymptotics, local uniqueness results, trace formulas, and Borg-type theorems for Dirac operators, *Trans. Amer. Math. Soc.* **354** (2002), no. 9, 3475–3534.
- [8] V. B. Daskalov and E. Kh. Khristov, Explicit formulae for the inverse problem for the regular Dirac operator, *Inverse Problems* **16** (2000), no. 1, 247–258.
- [9] T. T. Džabiev, The inverse problem for the Dirac equation with a singularity, *Akad. Nauk Azerbaïdžan. SSR Dokl.* **22** (1966), no. 11, 8–12 (Russian).
- [10] L. D. Faddeev, The inverse problem in the quantum theory of scattering, *Uspekhi Mat. Nauk* **14** (1959), pp. 57–119 (Russian); Engl. transl.: *J. Math. Phys.* **4** (1963), 72–104.
- [11] L. V. Fam, The inverse scattering problem for a system of Dirac equations on the whole axis, *Ukrain. Mat. Zh.* **24** (1972), 666–674 (Russian).
- [12] I. S. Frolov, An inverse scattering problem for the Dirac system on the entire axis, *Dokl. Akad. Nauk SSSR* **207** (1972), 44–47 (Russian).
- [13] M. G. Gasymov, An inverse problem of scattering theory for a system of Dirac equations of order $2n$, *Trudy Moskov. Mat. Ob-va* **19** (1968), 41–112 (Russian).
- [14] M. G. Gasymov and T. T. Džabiev, Solution of the inverse problem by two spectra for the Dirac equation on a finite interval, *Akad. Nauk Azerbaïdžan. SSR Dokl.* **22** (1966), no. 7, 3–6 (Russian).
- [15] M. G. Gasymov and B. M. Levitan, Determination of the Dirac system from the scattering phase, *Dokl. Akad. Nauk SSSR* **167** (1966), 1219–1222 (Russian).
- [16] M. G. Gasymov and B. M. Levitan, The inverse problem for the Dirac system, *Dokl. Akad. Nauk SSSR* **167** (1966), 967–970 (Russian).
- [17] F. Gesztesy, A. Kiselev, and K. A. Makarov, Uniqueness results for matrix-valued Schrödinger, Jacobi, and Dirac-type operators, *Math. Nachr.* **239/240** (2002), 103–145.
- [18] I. Gohberg, S. Goldberg, and M. Kaashoek, *Classes of linear operators. Vol. II*, Oper. Theory Adv. Appl. **63**, Birkhäuser Verlag, Basel, 1993.
- [19] I. Gohberg and M. Krein, *Introduction to the Theory of Linear Non-selfadjoint Operators in Hilbert Space*, Nauka Publ., Moscow, 1965 (Russian); Engl. transl.: Amer. Math. Soc. Transl. Math. Monographs **18**, Amer. Math. Soc., Providence, RI, 1969.
- [20] I. Gohberg and M. Krein, *Theory of Volterra Operators in Hilbert Space and its Applications*, Nauka Publ., Moscow, 1967 (Russian); Engl. transl.: Amer. Math. Soc. Transl. Math. Monographs **24**, Amer. Math. Soc., Providence, RI, 1970.
- [21] B. Grébert, Inverse scattering for the Dirac operator on the real line, *Inverse Problems* **8** (1992), no. 5, 787–807.
- [22] P. R. Halmos and V. S. Sunder, *Bounded Integral Operators on L^2 Spaces*, Springer-Verlag, Berlin-Heidelberg-New York, 1978.

- [23] D. B. Hinton, A. K. Jordan, M. Klaus, and J. K. Shaw, Inverse scattering on the line for a Dirac system, *J. Math. Phys.* **32** (1991), no. 11, 3015–3030.
- [24] M. Horváth, On a theorem of Ambarzumian, *Proc. Roy. Soc. Edinburgh Sect. A* **131** (2001), no. 4, 899–907.
- [25] M. Horváth, On the inverse spectral theory of Schrödinger and Dirac operators, *Trans. Amer. Math. Soc.* **353** (2001), no. 10, 4155–4171.
- [26] R. O. Hryniv and Ya. V. Mykytyuk, Inverse spectral problems for Sturm–Liouville operators with singular potentials, II. Reconstruction by two spectra, in *Functional Analysis and its Applications*, V. Kadets and W. Żelazko, eds., North-Holland Mathematics Studies, **197**, 97–114, North-Holland Publishing Co., Amsterdam, 2004.
- [27] R. Hryniv and Ya. Mykytyuk, On zeros of some entire functions, *arXiv preprint* <http://arxiv.org/abs/math.CV/0410420>.
- [28] Yi. Katznelson, *An Introduction to Harmonic Analysis*, 3rd edition, Cambridge University Press, Cambridge, 2004.
- [29] A. B. Khasanov and A. M. Ibragimov, On an inverse problem for the Dirac operator with periodic potential, *Uzbek. Mat. Zh.* **2001** no. 3-4, 48–55 (Russian).
- [30] M. Kiss, An n -dimensional Ambarzumian type theorem for Dirac operators, *Inverse Problems* **20** (2004), 1593–1597.
- [31] E. L. Korotyaev, Inverse problems for the Hill and Dirac operators, *Dokl. Akad. Nauk* **365** (1999), no. 6, 730–733 (Russian).
- [32] M. G. Krein, On integral equations generating differential equations of the second order, *Dokl. Akad. Nauk SSSR* **97** (1954), 21–24 (Russian).
- [33] M. Lesch and M. M. Malamud, The inverse spectral problem for first order systems on the half line, in *Differential operators and related topics, Vol. I (Odessa, 1997)*, 199–238, Oper. Theory Adv. Appl. **117**, Birkhäuser, Basel, 2000.
- [34] B. Ya. Levin, *Lectures on Entire Functions*, Amer. Math. Soc. Transl. Math. Monographs **150**, Amer. Math. Soc., Providence, RI, 1996.
- [35] N. Levinson, *Gap and Density Theorems*, Amer. Math. Soc. Colloquium Publ. **26**, Amer. Math. Soc., New York, 1940.
- [36] B. M. Levitan and I. S. Sargsjan, *Sturm–Liouville and Dirac operators*, Nauka Publ., Moscow, 1988 (Russian); Engl. transl.: Kluwer Acad. Publ., Dordrecht, 1991.
- [37] F. G. Maksudov and S. G. Veliev, An inverse scattering problem for the nonselfadjoint Dirac operator on the whole axis, *Dokl. Akad. Nauk SSSR* **225** (1975), no. 6, 1263–1266 (Russian).
- [38] M. M. Malamud, On Borg-type theorems for first-order systems on a finite interval, *Funktsional. Anal. i Prilozhen.* **33** (1999), no. 1, 75–80 (Russian); Engl. transl.: *Funct. Anal. Appl.* **33** (1999), no. 1, 64–68.
- [39] M. M. Malamud, Questions of uniqueness in inverse problems for systems of differential equations on a finite interval, *Tr. Mosk. Mat. Ob-va* **60** (1999), 199–258 (Russian); Engl. transl.: *Trans. Moscow Math. Soc.* **1999**, 173–224.
- [40] S. G. Mamedov, The inverse boundary value problem on a finite interval for Dirac’s system of equations, *Azerbaïdžan. Gos. Univ. Učen. Zap. Ser. Fiz-Mat. Nauk*, 1975, no. 5, 61–67 (Russian).
- [41] V. A. Marchenko, *Sturm–Liouville Operators and Their Applications*, Naukova Dumka Publ., Kiev, 1977 (Russian); Engl. transl.: Birkhäuser Verlag, Basel, 1986.
- [42] K. Mochizuki and I. Trooshin, Inverse problem for interior spectral data of the Dirac operator on a finite interval, *Publ. Res. Inst. Math. Sci.* **38** (2002), no. 2, 387–395.
- [43] Ya. V. Mykytyuk, Factorization of Fredholm operators, *Mat. Stud.* **20** (2003), no. 2, 185–199 (Ukrainian).
- [44] Ya. V. Mykytyuk, Factorization of Fredholm operators in operator algebras, *Mat. Stud.* **21** (2004), no. 1, 87–97 (Ukrainian).
- [45] I. M. Nabiev, Solution of a class of inverse problems for the Dirac operator, *Trans. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci.* **21** (2001), no. 1, 146–157 (Russian).
- [46] M. A. Naïmark, *Normed Algebras*, 2nd revised ed., Nauka Publ., Moscow, 1968 (Russian); Engl. transl.: 3rd ed., Wolters-Noordhoff Series of Monographs and Textbooks on Pure and Applied Mathematics, Wolters-Noordhoff Publishing, Groningen, 1972.

- [47] L. P. Nizhnik, An inverse problem of nonstationary scattering for the Dirac equations, *Ukrain. Mat. Zh.* **24** (1972), 110–113 (Russian).
- [48] L. P. Nizhnik, M. D. Pochinaiko, and V. G. Tarasov, The inverse scattering problem for a system of Dirac equations in characteristic variables, in *The Spectral Theory of Operators in Problems of Mathematical Physics*, 72–93, Akad. Nauk Ukrain. SSR, Inst. Mat., Kiev, 1983 (Russian).
- [49] A. Sakhnovich, Dirac type and canonical systems: spectral and Weyl-Titchmarsh matrix functions, direct and inverse problems, *Inverse Problems* **18** (2002), no. 2, 331–348.
- [50] A. Sakhnovich, Dirac type system on the axis: explicit formulae for matrix potentials with singularities and soliton-positon interactions, *Inverse Problems* **19** (2003), no. 4, 845–854.
- [51] I. S. Sargsjan, A uniqueness theorem for the solution of the inverse problem for a one-dimensional Dirac system, in *Some Boundary Value Problems of Ordinary Differential Equations*, 3–13, Univ. Družby Narodov, Moscow, 1970 (Russian).
- [52] A. M. Savchuk and A. A. Shkalikov, The Sturm-Liouville operators with distributional potential, *Trudy Mosk. Matem. Ob-va*, **64** (2003), 159–212 (Russian); Engl. transl.: *Trans. Moscow Math. Soc.* **2003**, 143–192.
- [53] D. G. Shepel'skiĭ, An inverse spectral problem for a Dirac-type operator with "sewing", in *Dynamical Systems and Complex Analysis*, 104–112, "Naukova Dumka", Kiev, 1992 (Russian).
- [54] B. Thaller, *The Dirac Equation*, Springer, Berlin, 1992.
- [55] B. A. Watson, Inverse spectral problems for weighted Dirac systems, *Inverse Problems* **15** (1999), no. 3, 793–805.
- [56] V. E. Zakharov and A. B. Shabat, Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media, *Zh. Èksper. Teoret. Fiz.* **61** (1971), no. 1, 118–134 (Russian); Engl. transl.: *Soviet Physics JETP* **34** (1972), no. 1, 62–69.

(S.A.) INSTITUT FÜR ANGEWANDTE MATHEMATIK, UNIVERSITÄT BONN, WEGELERSTR. 6, D-53115, BONN, GERMANY; SFB 611, BONN, GERMANY; BiBoS, BIELEFELD, GERMANY; IZKS; CERFIM, LOCARNO, SWITZERLAND; AND ACCADEMIA DI ARCHITETTURA, MENDRISIO, SWITZERLAND

E-mail address: `albeverio@uni-bonn.de`

(R.H. and Ya.M.) INSTITUTE FOR APPLIED PROBLEMS OF MECHANICS AND MATHEMATICS, 3B NAUKOVA ST., 79601 LVIV, UKRAINE AND LVIV NATIONAL UNIVERSITY, 1 UNIVERSYTETSKA ST., 79602 LVIV, UKRAINE

E-mail address: `rhryniv@iapmm.lviv.ua` and `yamykytyuk@yahoo.com`